

CLASSIFICATION OF ABELIAN COMPLEX STRUCTURES ON 6-DIMENSIONAL LIE ALGEBRAS

A. ANDRADA, M.L. BARBERIS, AND I.G. DOTTI

ABSTRACT. We classify the 6-dimensional Lie algebras that can be endowed with an abelian complex structure and parameterize, on each of these algebras, the space of such structures up to holomorphic isomorphism.

1. INTRODUCTION

Let $\mathfrak{g}$ be a Lie algebra, J an endomorphism of $\mathfrak{g}$ such that $J^2 = -I$, and let $\mathfrak{g}^{1,0}$ be the i -eigenspace of J in $\mathfrak{g}^{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$. When $\mathfrak{g}^{1,0}$ is a complex subalgebra we say that J is integrable, when $\mathfrak{g}^{1,0}$ is abelian we say that J is abelian and when $\mathfrak{g}^{1,0}$ is a complex ideal we say that J is bi-invariant. We note that a complex structure on a Lie algebra cannot be both abelian and bi-invariant, unless the Lie bracket is trivial. If G is a connected Lie group with Lie algebra $\mathfrak{g}$, by left translating J one obtains a complex manifold (G, J) such that left multiplication is holomorphic and, in the bi-invariant case, also right multiplication is holomorphic, which implies that (G, J) is a complex Lie group.

Our concern here will be the case when J is abelian. In this case the Lie algebra has abelian commutator, thus, it is 2-step solvable (see [16]). However, its nilradical need not be abelian (see Remark 4.7). Abelian complex structures have interesting applications in hyper-Kähler with torsion geometry (see [6]). It has been shown in [8] that the Dolbeault cohomology of a nilmanifold with an abelian complex structure can be computed algebraically. Also, deformations of abelian complex structures on nilmanifolds have been studied in [9].

Of importance, when studying complex structures on a Lie algebra $\mathfrak{g}$, is the ideal $\mathfrak{g}'_J := \mathfrak{g}' + J\mathfrak{g}'$ constructed from algebraic and complex data. We will say that the complex structure J is proper when $\mathfrak{g}'_J$ is properly contained in $\mathfrak{g}$. Any complex structure on a nilpotent Lie algebra is proper [18]. The classification of nilpotent 6-dimensional Lie algebras carrying complex structures has been given in [18]; when the complex structure is abelian the characterization was obtained in [11].

There is only one 2-dimensional non-abelian Lie algebra, the Lie algebra of the affine motion group of $\mathbb{R}$, denoted by $\mathfrak{aff}(\mathbb{R})$. It carries a unique complex structure, up to equivalence, which turns out to be abelian. The 4-dimensional Lie algebras admitting abelian complex structures were classified in [19]. Each of these Lie algebras, with the exception of $\mathfrak{aff}(\mathbb{C})$, the realification of the Lie algebra of the affine motion group of $\mathbb{C}$, has a unique abelian complex structure up to equivalence. On $\mathfrak{aff}(\mathbb{C})$ there is a two-sphere of abelian complex structures, but only two equivalence classes distinguished by J being proper or not. Furthermore, $\mathfrak{aff}(\mathbb{C})$ is equipped with a natural bi-invariant complex structure. In dimension 6 it turns out that, as a consequence of our results, some of the Lie algebras equipped with abelian complex structures are of the form $\mathfrak{aff}(A)$ where A is a 3-dimensional commutative associative algebra.

1991 *Mathematics Subject Classification.* Primary 17B30; Secondary 53C15.

Key words and phrases. Abelian complex structures, nilpotent and solvable Lie algebras.

In this article we obtain the classification of the 6-dimensional Lie algebras that can be endowed with an abelian complex structure. Moreover, on each of the resulting Lie algebras, we parameterize the space of such structures up to holomorphic isomorphism. It turns out that there are three nilpotent Lie algebras carrying curves of non-equivalent structures, while for the remaining Lie algebras the moduli space is finite.

The outline of the paper is as follows. In §2 we give the basic definitions, recall known results and study some properties of the Lie algebra $\mathfrak{aff}(\mathbb{C})$. In addition, we parameterize the space of equivalence classes of complex structures on a Lie algebra with 1-dimensional commutator (see Proposition 2.2). In §3 we perform the classification in the 6-dimensional nilpotent case. This is done in two steps. Firstly, we obtain the classification of the nilpotent Lie algebras admitting such a structure (see Theorem 3.2), recovering, by a different approach, the results obtained in [11, 18]. In the second step, we parameterize, up to equivalence, the space of abelian complex structures on each of these algebras (see Theorem 3.5). As a consequence, we obtain three nilpotent Lie algebras carrying curves of non-equivalent abelian complex structures, namely, $\mathfrak{h}_3 \times \mathfrak{h}_3$, $\mathfrak{h}_3(\mathbb{C})$ and a 3-step nilpotent Lie algebra, where $\mathfrak{h}_3$ is the 3-dimensional real Heisenberg Lie algebra and $\mathfrak{h}_3(\mathbb{C})$ is the realification of the 3-dimensional complex Heisenberg Lie algebra.

In §4 we determine the 6-dimensional non-nilpotent Lie algebras $\mathfrak{s}$ with an abelian complex structure, studying in Theorems 4.1, 4.2 and 4.4 the case when such a structure is proper. The Lie algebras, a finite number, appearing in Theorems 4.1 and 4.2 are decomposable as a direct product of lower dimensional Lie algebras with the corresponding abelian complex structure preserving this decomposition. In Theorem 4.4, when considering the case $\mathfrak{s}'_J = \mathbb{R}^4$, we obtain a 2-parameter family of solvable Lie algebras carrying both an abelian and a bi-invariant complex structure. Moreover, as a consequence of this result and Theorems 4.1 and 4.2, it follows that the Lie algebra $\mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ has exactly three abelian complex structures, up to equivalence, distinguished by the ideal $\mathfrak{s}'_J$. Finally, in Theorem 4.6 we consider the case when J is not proper and obtain five isomorphism classes of Lie algebras, all of which are of the form $\mathfrak{aff}(A)$ where A is a commutative associative algebra with identity.

2. PRELIMINARIES

2.1. Complex structures on Lie algebras. A complex structure on a real Lie algebra $\mathfrak{g}$ is an endomorphism J of $\mathfrak{g}$ satisfying $J^2 = -I$ and such that

$$(1) \quad J[x, y] - [Jx, y] - [x, Jy] - J[Jx, Jy] = 0, \quad \forall x, y \in \mathfrak{g}.$$

It is well known that (1) holds if and only if $\mathfrak{g}^{1,0}$, the i -eigenspace of J , is a complex subalgebra of $\mathfrak{g}^{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$. A pair $(\mathfrak{g}, J)$ will denote a Lie algebra $\mathfrak{g}$ equipped with a complex structure J .

A complex structure J on $\mathfrak{g}$ is called *bi-invariant* if the following condition holds:

$$(2) \quad J[x, y] = [Jx, y], \quad x, y \in \mathfrak{g}.$$

In this case, $(\mathfrak{g}, J)$ is a complex Lie algebra, that is, $\mathfrak{g}$ is turned into a $\mathbb{C}$ -vector space by setting

$$(a + ib)x = ax + bJx, \quad x \in \mathfrak{g}, \quad a, b \in \mathbb{R},$$

and condition (2) ensures that the Lie bracket is now $\mathbb{C}$ -bilinear.

Two complex structures J_1 and J_2 on $\mathfrak{g}$ are said to be *equivalent* if there exists an automorphism α of $\mathfrak{g}$ satisfying $J_2 \alpha = \alpha J_1$. Two pairs $(\mathfrak{g}_1, J_1)$ and $(\mathfrak{g}_2, J_2)$ are *holomorphically isomorphic* if there exists a Lie algebra isomorphism $\alpha : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ satisfying $J_2 \alpha = \alpha J_1$.

Consider a Lie algebra $\mathfrak{g}$ equipped with a complex structure J . We denote with $\mathfrak{g}'$ the commutator ideal $[\mathfrak{g}, \mathfrak{g}]$ of $\mathfrak{g}$ and with $\mathfrak{g}'_J$ the J -stable ideal of $\mathfrak{g}$ defined by $\mathfrak{g}'_J := \mathfrak{g}' + J\mathfrak{g}'$. We will say that J is *proper* if $\mathfrak{g}'_J \subsetneq \mathfrak{g}$. We note that if $(\mathfrak{g}_1, J_1)$ and $(\mathfrak{g}_2, J_2)$ are holomorphically isomorphic, then J_1 is proper if and only if J_2 is.

The next proposition provides sufficient conditions for a complex structure to be proper. We recall that a Lie algebra is called *perfect* if it coincides with its commutator ideal.

Proposition 2.1. (1) *Every complex structure on a nilpotent Lie algebra is proper.*

(2) *Every bi-invariant complex structure on a non-perfect Lie algebra is proper.*

Proof. The first statement was proved in [18, Theorem 1.3] (see also Lemma 2.4 below). The second assertion follows since (2) implies that the commutator ideal is stable by any bi-invariant complex structure. $\square$

2.2. Abelian complex structures on Lie algebras. An *abelian* complex structure on $\mathfrak{g}$ is an endomorphism J of $\mathfrak{g}$ satisfying

$$(3) \quad J^2 = -I, \quad [Jx, Jy] = [x, y], \quad \forall x, y \in \mathfrak{g}.$$

It follows that (3) is a particular case of (1); moreover, condition (3) is equivalent to $\mathfrak{g}^{1,0}$ being abelian. These structures were first considered in [3]. A class of Lie algebras carrying such structures appears in the next result.

Let $\mathfrak{h}_{2n+1} = \text{span}\{e_1, \dots, e_{2n}, z_0\}$ denote the $(2n+1)$ -dimensional Heisenberg algebra with Lie bracket $[e_{2i-1}, e_{2i}] = z_0$, $1 \leq i \leq n$, and $\mathfrak{aff}(\mathbb{R})$ the Lie algebra of the group of affine motions of $\mathbb{R}$.

Proposition 2.2. *If $\mathfrak{g}$ is an even dimensional real Lie algebra with 1-dimensional commutator $\mathfrak{g}'$, then:*

- (1) $\mathfrak{g}$ is isomorphic to either $\mathfrak{h}_{2n+1} \times \mathbb{R}^{2k+1}$ or $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^{2k}$;
- (2) All these Lie algebras carry abelian complex structures and every complex structure on $\mathfrak{g}$ is abelian;
- (3) There are $\left\lfloor \frac{n}{2} \right\rfloor + 1$ equivalence classes of complex structures on $\mathfrak{h}_{2n+1} \times \mathbb{R}^{2k+1}$;
- (4) There is a unique complex structure on $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^{2k}$ up to equivalence.

Proof. (1) follows from the characterization of the Lie algebras with one dimensional commutator (see for example [4, Theorem 4.1]).

A proof of (2) can be found in [5, Examples 3.3 and Proposition 3.4].

In order to prove (3), we recall that according to [17, Proposition 3.6] the complex structures on $\mathfrak{h}_{2n+1} \times \mathbb{R}^{2k+1}$ are given by:

$$(4) \quad J e_{2i-1} = \pm e_{2i}, \quad J z_j = z_{j+1}, \quad 1 \leq i \leq n, \quad 0 \leq j \leq k,$$

where $\{e_1, \dots, e_{2n}, z_0\}$ is a basis of $\mathfrak{h}_{2n+1}$ such that $[e_{2i-1}, e_{2i}] = z_0$ and $\{z_1, \dots, z_{2k+1}\}$ is a basis of $\mathbb{R}^{2k+1}$. Any two complex structures with the same number of $-$ signs are equivalent, by permuting the elements of the basis. Let J_r , with $0 \leq r \leq n$, be any complex structure as in (4) such that r is the number of $-$ signs. By a suitable permutation of the basis, one can show that J_r is equivalent to J_{n-r} , thus we may assume $0 \leq r \leq \left\lfloor \frac{n}{2} \right\rfloor$. Furthermore, for $0 \leq r, s \leq \left\lfloor \frac{n}{2} \right\rfloor$, if J_r is equivalent to J_s , by counting the multiplicities of the eigenvalues of J_r and J_s on $\mathfrak{g} \otimes \mathbb{C}$, we must have $r = s$ and (3) follows.¹

¹ The classification for $n = 1$, $k = 0$, was given in [19] and the case $n = 2$, $k = 0$ was carried out in [20].

The proof of (4) follows by observing that given a complex structure J on $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^{2k}$, there exists a J -stable complementary ideal $\mathfrak{s}$ of $\mathbb{R}^{2k}$ isomorphic to $\mathfrak{aff}(\mathbb{R})$. It is clear that $\mathfrak{aff}(\mathbb{R})$ admits a unique complex structure up to equivalence, and the assertion follows. $\square$

We include some properties of abelian complex structures in the following lemma, whose proof is straightforward.

Lemma 2.3. *Let J be an abelian complex structure on a Lie algebra $\mathfrak{g}$. Then:*

- (i) *the center $\mathfrak{z}(\mathfrak{g})$ of $\mathfrak{g}$ is J -stable;*
- (ii) *for any $x \in \mathfrak{g}$, $\text{ad}_{Jx} = -\text{ad}_x J$; in particular, the family of inner derivations of $\mathfrak{g}$ is stable under right multiplication by J ;*
- (iii) *for any J -stable ideal $\mathfrak{u}$ of $\mathfrak{g}$, the kernel of the map $\mathfrak{g} \rightarrow \text{End}(\mathfrak{u})$, $x \mapsto \text{ad}_x|_{\mathfrak{u}}$, is J -stable.*

Some known obstructions to the existence of abelian complex structures are stated below.

- (O1) Let $\mathfrak{g}$ be a real Lie algebra admitting an abelian complex structure. Then $\mathfrak{g}$ is 2-step solvable [16].
- (O2) Let $\mathfrak{g}$ be a solvable Lie algebra such that $\mathfrak{g}'$ has codimension 1 in $\mathfrak{g}$ and $\dim \mathfrak{g} > 2$. Then $\mathfrak{g}$ does not admit abelian complex structures [5].

Another obstruction in the case of nilpotent Lie algebras is given in the next lemma. Recall that the descending central series of $\mathfrak{g}$ is defined by $\mathfrak{g}^0 = \mathfrak{g}$, $\mathfrak{g}^i = [\mathfrak{g}, \mathfrak{g}^{i-1}]$, $i \geq 1$.

Lemma 2.4. *Let $\mathfrak{g}$ be a k -step nilpotent Lie algebra with an abelian complex structure J and set $\mathfrak{g}_J^i := \mathfrak{g}^i + J\mathfrak{g}^i$. Then $\mathfrak{g}_J^i \subsetneq \mathfrak{g}_J^{i-1}$ for all $i \leq k$. In particular, if $\dim \mathfrak{g} = 2m$, $\mathfrak{g}$ is at most m -step nilpotent.*

Proof. Clearly, $\{0\} = \mathfrak{g}_J^k \subsetneq \mathfrak{g}_J^{k-1}$. Assume that $\mathfrak{g}_J^i = \mathfrak{g}_J^{i-1}$ for some $i < k$. Then, any $x \in \mathfrak{g}^{i-1}$ can be written as $x = y + Jw$, $y, w \in \mathfrak{g}^i$. The fact that J is abelian implies that $[\mathfrak{g}, x] \subset \mathfrak{g}^{i+1}$ for all $x \in \mathfrak{g}^{i-1}$, that is, $\mathfrak{g}^i \subset \mathfrak{g}^{i+1}$, contradicting that $\mathfrak{g}$ is k -step nilpotent. $\square$

2.3. Dimension less than or equal to four. There are only two 2-dimensional Lie algebras: $\mathbb{R}^2$ and $\mathfrak{aff}(\mathbb{R})$. Both carry a unique abelian complex structure, up to equivalence.

Let $\mathfrak{g}$ be a 4-dimensional solvable Lie algebra with basis $\{e_1, e_2, e_3, e_4\}$. According to [19], if $\mathfrak{g}$ admits an abelian complex structure, it is isomorphic to one and only one of the following:

$$\begin{aligned} \mathfrak{g}_1 &= \mathbb{R}^4, \\ \mathfrak{g}_2 &= \mathfrak{h}_3 \times \mathbb{R}, [e_1, e_2] = e_3, \\ \mathfrak{g}_3 &= \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2, [e_1, e_2] = e_2 \\ \mathfrak{g}_4 &= \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}), [e_1, e_2] = e_2, [e_3, e_4] = e_4 \\ \mathfrak{g}_5 &= \mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2, [e_1, e_2] = e_2, [e_1, e_4] = e_4, [e_2, e_3] = e_4. \\ \mathfrak{g}_6 &= \mathfrak{aff}(\mathbb{C}), [e_1, e_3] = e_3, [e_1, e_4] = e_4, [e_2, e_3] = e_4, [e_2, e_4] = -e_3, \end{aligned}$$

where $\mathfrak{h}_3$ is the 3-dimensional Heisenberg algebra and $\mathfrak{aff}(\mathbb{C})$ is the realification of the Lie algebra of the group of affine motions of the complex line.

Theorem 2.5 ([19]). *Let $\mathfrak{g}$ be a 4-dimensional solvable Lie algebra with an abelian complex structure J . Then $(\mathfrak{g}, J)$ is holomorphically isomorphic to one and only one of the following:*

$$\begin{aligned} \mathfrak{g}_1 : J e_1 &= e_2, J e_3 = e_4, \\ \mathfrak{g}_2 : J e_1 &= e_2, J e_3 = e_4, \end{aligned}$$

$$\begin{aligned}
\mathfrak{g}_3 : J e_1 &= e_2, J e_3 = e_4, \\
\mathfrak{g}_4 : J e_1 &= e_2, J e_3 = e_4, \\
\mathfrak{g}_5 : J e_1 &= e_2, J e_3 = -e_4, \\
\mathfrak{g}_6 : J_1 e_1 &= -e_2, J_1 e_3 = e_4, \\
\mathfrak{g}_6 : J_2 e_1 &= e_3, J_2 e_2 = e_4.
\end{aligned}$$

We include below a proof of the fact that, up to equivalence, $\mathfrak{aff}(\mathbb{C})$ has two abelian complex structures (see also [19]). We recall that $\mathfrak{aff}(\mathbb{C})$ has a basis $\{e_1, e_2, e_3, e_4\}$ with the following Lie bracket

$$(5) \quad [e_1, e_3] = e_3, \quad [e_1, e_4] = e_4, \quad [e_2, e_3] = e_4, \quad [e_2, e_4] = -e_3,$$

and with automorphism group given by (see for instance [7]):

$$(6) \quad \text{Aut}(\mathfrak{aff}(\mathbb{C})) = \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \epsilon & 0 & 0 \\ c & -\epsilon d & a & -\epsilon b \\ d & \epsilon c & b & \epsilon a \end{pmatrix} : a, b, c, d, \in \mathbb{R}, \epsilon = \pm 1 \right\}.$$

Proposition 2.6. *Any abelian complex structure on the Lie algebra $\mathfrak{aff}(\mathbb{C})$ is equivalent to either J_1 or J_2 , where*

$$\begin{aligned}
J_1 e_1 &= -e_2, & J_1 e_3 &= e_4, \\
J_2 e_1 &= e_3, & J_2 e_2 &= e_4.
\end{aligned}$$

Moreover, J_1 is proper and J_2 is not, hence they are not equivalent.

Proof. Let us assume that J is an abelian complex structure on $\mathfrak{g} = \mathfrak{aff}(\mathbb{C})$. Since $\dim \mathfrak{g}' = 2$, there are two possibilities for $\mathfrak{g}'$:

- $\mathfrak{g}'_J = \mathfrak{g}'$. In this case, $\mathfrak{g}'$ is a J -stable ideal of $\mathfrak{aff}(\mathbb{C})$. Let

$$J = \begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$$

be the matrix representation of J with respect to the ordered basis $\{e_1, e_2, e_3, e_4\}$. Since $J^2 = -I$, we obtain that $A^2 = -I$ and $C^2 = -I$, and therefore J can be written as

$$J = \begin{pmatrix} \alpha & \gamma & 0 & 0 \\ \beta & -\alpha & 0 & 0 \\ a & c & e & g \\ b & d & f & -e \end{pmatrix},$$

with

$$(7) \quad \alpha^2 + \beta\gamma = -1, \quad e^2 + fg = -1,$$

and certain constraints on a, b, c, d . From the fact that J is abelian, together with (7), we obtain that

$$\alpha = e = 0, \quad \beta = \pm 1, \quad f = -g = \gamma = -\frac{1}{\beta}.$$

Using again that $J^2 = -I$, we arrive at $c = -b$, $d = a$, thus, $J = J_{(a,b)}^+$ or $J = J_{(a,b)}^-$, where

$$J_{(a,b)}^+ := \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ a & -b & 0 & -1 \\ b & a & 1 & 0 \end{pmatrix}, \quad J_{(a,b)}^- := \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ a & -b & 0 & 1 \\ b & a & -1 & 0 \end{pmatrix}.$$

Note that $J_{(0,0)}^+ = J_1$. Consider now the automorphisms ϕ and ψ of $\mathfrak{aff}(\mathbb{C})$ given by

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ \frac{b}{2} & \frac{a}{2} & 1 & 0 \\ \frac{a}{2} & -\frac{b}{2} & 0 & -1 \end{pmatrix}, \quad \psi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -\frac{b}{2} & -\frac{a}{2} & 1 & 0 \\ \frac{a}{2} & -\frac{b}{2} & 0 & 1 \end{pmatrix}.$$

It is easy to see that $\phi J_{(a,b)}^+ = J_1 \phi$ and $\psi J_{(a,b)}^- = J_1 \psi$, so that all the abelian complex structures $J_{(a,b)}^+$ and $J_{(a,b)}^-$ are equivalent to J_1 . This completes this case.

• $\mathfrak{g}'_J = \mathfrak{g}$. In this case $\mathfrak{g} = \mathfrak{g}' \oplus J\mathfrak{g}'$. Let

$$f_1 = Je_3 = ae_1 + be_2 + v, \quad f_2 = Je_4 = ce_1 + de_2 + w,$$

$a, b, c, d \in \mathbb{R}$, $v, w \in \text{span}\{e_3, e_4\}$. Since J is abelian, $[f_1, f_2] = 0$ and $[f_2, e_3] = [f_1, e_4]$, hence $c = -b$, $d = a$. Therefore, we have

$$\text{ad}_{f_1}|_{\mathfrak{g}'} = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}, \quad \text{ad}_{f_2}|_{\mathfrak{g}'} = \begin{pmatrix} -b & -a \\ a & -b \end{pmatrix}.$$

Observe that $a^2 + b^2 \neq 0$, since otherwise $\mathfrak{g}$ would be abelian. Setting

$$\tilde{f}_1 = \frac{1}{a^2 + b^2}(af_1 - bf_2), \quad \tilde{f}_2 = \frac{1}{a^2 + b^2}(bf_1 + af_2),$$

we obtain

$$\text{ad}_{\tilde{f}_1}|_{\mathfrak{g}'} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \text{ad}_{\tilde{f}_2}|_{\mathfrak{g}'} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The expression of J with respect to the basis $\{\tilde{f}_1, \tilde{f}_2, e_3, e_4\}$ is given by

$$J = J_{(a,b)} = \begin{pmatrix} & B \\ -B^{-1} & \end{pmatrix}, \quad \text{where } B = \begin{pmatrix} -a & b \\ -b & -a \end{pmatrix}.$$

The structure constants of $\mathfrak{g}$ with respect to the new basis $\{\tilde{f}_1, \tilde{f}_2, e_3, e_4\}$ are given by (5). If ϕ is the following automorphism of $\mathfrak{aff}(\mathbb{C})$:

$$\phi = \begin{pmatrix} 1 & 0 & & \\ 0 & 1 & & \\ & & -a & b \\ & & -b & -a \end{pmatrix},$$

it turns out that $\phi J_{(a,b)} = J_{(1,0)} \phi$, that is, $J_{(a,b)}$ is equivalent to $J_{(1,0)} = J_2$, and the proposition follows. $\square$

Given $x = (x_1, x_2, x_3) \in S^2$, let $J_x := x_1 J_1 + x_2 J_2 + x_3 J_1 J_2$ with J_1 and J_2 as above. Since $J_1 J_2 = -J_2 J_1$, J_x is an abelian complex structure on $\mathfrak{aff}(\mathbb{C})$ and Proposition 2.6 implies the following result:

Corollary 2.7. *There is a two-sphere J_x , $x \in S^2$, of abelian complex structures on $\mathfrak{aff}(\mathbb{C})$. Moreover, J_x is equivalent to J_1 if and only if $x = (1, 0, 0)$ or $x = (-1, 0, 0)$. All the other abelian complex structures in this sphere are equivalent to J_2 .*

Remark 2.8. It was shown in [2] that $\mathfrak{aff}(\mathbb{C})$ is the only 4-dimensional Lie algebra carrying an abelian hypercomplex structure, that is, a pair of anticommuting abelian complex structures.

Remark 2.9. We point out that $\mathfrak{aff}(\mathbb{C})$ admits a bi-invariant complex structure J , namely,

$$Je_1 = e_2, \quad Je_3 = e_4.$$

Moreover, any bi-invariant complex structure is given by $\pm J$ and J is equivalent to $-J$ via an automorphism as in (6) with $\epsilon = -1$. Therefore, $\mathfrak{aff}(\mathbb{C})$ has a unique bi-invariant complex structure up to equivalence.

The next characterization of $(\mathfrak{aff}(\mathbb{C}), J_1)$ will be frequently used in §4.

Lemma 2.10. *Let $\mathfrak{g}$ be a 4-dimensional Lie algebra spanned by $\{f_1, \dots, f_4\}$ with Lie bracket defined by:*

$$\begin{aligned} [f_1, f_3] &= [f_2, f_4] = cf_3 + df_4, & [f_1, f_4] &= -[f_2, f_3] = -df_3 + cf_4, \\ [f_1, f_2] &= xf_3 + yf_4, \end{aligned}$$

with $c^2 + d^2 \neq 0$, and abelian complex structure J given by

$$Jf_1 = f_2, \quad Jf_3 = f_4.$$

Then $(\mathfrak{g}, J)$ is holomorphically isomorphic to $(\mathfrak{aff}(\mathbb{C}), J_1)$.

Proof. Consider first

$$\tilde{f}_1 = \frac{cf_1 + df_2}{c^2 + d^2}, \quad \tilde{f}_2 = J\tilde{f}_1 = \frac{-df_1 + cf_2}{c^2 + d^2}.$$

In the new basis $\{\tilde{f}_1, \tilde{f}_2, f_3, f_4\}$, the Lie bracket of $\mathfrak{g}$ becomes

$$[\tilde{f}_1, \tilde{f}_2] = x'f_3 + y'f_4, \quad [\tilde{f}_1, f_3] = [\tilde{f}_2, f_4] = f_3, \quad [\tilde{f}_1, f_4] = -[\tilde{f}_2, f_3] = f_4.$$

Setting now

$$e_1 := \tilde{f}_1 - \frac{y'}{2}f_3 + \frac{x'}{2}f_4, \quad e_2 := -\tilde{f}_2 + \frac{x'}{2}f_3 + \frac{y'}{2}f_4, \quad e_3 := f_3, \quad e_4 := f_4,$$

we obtain a basis $\{e_1, \dots, e_4\}$ of $\mathfrak{g}$ such that the Lie bracket is given as in (5) and the complex structure J coincides with J_1 from Proposition 2.6. $\square$

3. DIMENSION SIX: THE NILPOTENT CASE

In this section we classify, up to holomorphic isomorphism, the 6-dimensional nilpotent non-abelian Lie algebras $\mathfrak{n}$ with abelian complex structures.

We begin by recalling from Proposition 2.1 that any complex structure J on a nilpotent Lie algebra $\mathfrak{n}$ is proper, that is, $\mathfrak{n}'_J = \mathfrak{n}' + J\mathfrak{n}'$ is a proper nilpotent ideal. In the 6-dimensional case, as a consequence of the following general result, it turns out that $\mathfrak{n}'_J$ is abelian.

Proposition 3.1. *If J is an abelian complex structure on a 6-dimensional solvable Lie algebra $\mathfrak{s}$ such that $\mathfrak{s}'_J$ is nilpotent, then $\mathfrak{s}'_J$ is abelian.*

Proof. If $\mathfrak{s}'_J = \mathfrak{s}$ then $\mathfrak{s}$ is nilpotent and Proposition 2.1 implies that $\mathfrak{s}'_J \subsetneq \mathfrak{s}$, a contradiction. Therefore, $\dim \mathfrak{s}'_J = 2$ or 4. If $\dim \mathfrak{s}'_J = 2$ then $\mathfrak{s}'_J$ is abelian.

We show next that if $\dim \mathfrak{s}'_J = 4$ and $\mathfrak{s}'_J$ is non-abelian nilpotent, then we get a contradiction. J induces by restriction an abelian complex structure on $\mathfrak{s}'_J$. The only 4-dimensional non-abelian nilpotent Lie algebra admitting abelian complex structures is $\mathfrak{h}_3 \times \mathbb{R}$ and this complex structure is unique up to equivalence (Theorem 2.5). Therefore, $\mathfrak{s}'_J \cong \mathfrak{h}_3 \times \mathbb{R}$, and there exists a basis $e_1, \dots, e_4$ of $\mathfrak{s}'_J$ such that $[e_1, e_2] = e_3$ and $Je_1 = e_2$, $Je_3 = e_4$. Let $\text{span}\{f_1, f_2\}$ be a complementary subspace of $\mathfrak{s}'_J$ such that $f_2 = Jf_1$. Setting $D = \text{ad}_{f_1}|_{\mathfrak{s}'_J}$, it follows from Lemma 2.3(ii) that $\text{ad}_{f_2}|_{\mathfrak{s}'_J} = -DJ$. The Lie bracket $[f_1, f_2]$ lies in $\mathfrak{s}' \subseteq \mathfrak{s}'_J$, hence, $[f_1, f_2] = ae_1 + be_2 + ce_3 + de_4$, with $a, b, c, d \in \mathbb{R}$.

Since D is a derivation of $\mathfrak{h}_3 \times \mathbb{R}$, it is of the form

$$D = \begin{pmatrix} A & 0 \\ B & C \end{pmatrix},$$

where A, B, C are 2×2 real matrices such that $c_{11} = \text{tr } A$ and $c_{21} = 0$. Moreover, DJ is also a derivation of $\mathfrak{h}_3 \times \mathbb{R}$, and therefore $c_{12} = a_{12} - a_{21}$ and $c_{22} = 0$. Observe that $\mathfrak{s}'_J$ is 2-step solvable and $\mathfrak{u} := \text{span}\{e_3, e_4\} \subset \mathfrak{s}'_J$ is an abelian ideal containing $(\mathfrak{s}'_J)'$, therefore the Jacobi identity implies²

$$C^2 J|_{\mathfrak{u}} = C(J|_{\mathfrak{u}})C,$$

and it follows that $C = 0$. Therefore, A takes the following form:

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & -a_{11} \end{pmatrix}.$$

Since $\mathfrak{s}'$ is abelian, we have that $\text{rank } A \leq 1$, hence $a_{11} = a_{12} = 0$, that is, $A = 0$. Finally, the Jacobi identity applied to f_1, f_2, e_j , $j = 1, 2$, implies that $a = b = 0$, that is, $[f_1, f_2] \in \mathfrak{u}$, hence $\mathfrak{s}'_J = \mathfrak{u}$, a contradiction. $\square$

In the remainder of this section we perform the classification. This will be done in two steps. In the first one, we obtain the classification of the Lie algebras admitting such a structure (see Theorem 3.2), recovering by a different approach the results obtained in [11, 18]. In the second step, we parameterize the space of abelian complex structures up to equivalence on each of these algebras (see Theorem 3.5).

Theorem 3.2. *Let $\mathfrak{n}$ be a 6-dimensional nilpotent Lie algebra with an abelian complex structure J . Then $\mathfrak{n}$ is isomorphic to one and only one of the following Lie algebras:*

$$\begin{aligned} \mathfrak{n}_1 &:= \mathfrak{h}_3 \times \mathbb{R}^3 : [e_1, e_2] = e_6, \\ \mathfrak{n}_2 &:= \mathfrak{h}_5 \times \mathbb{R} : [e_1, e_2] = e_6 = [e_3, e_4], \\ \mathfrak{n}_3 &:= \mathfrak{h}_3 \times \mathfrak{h}_3 : [e_1, e_2] = e_5, \quad [e_3, e_4] = e_6, \\ \mathfrak{n}_4 &:= \mathfrak{h}_3(\mathbb{C}) : [e_1, e_3] = -[e_2, e_4] = e_5, \quad [e_1, e_4] = [e_2, e_3] = e_6, \\ \mathfrak{n}_5 &: [e_1, e_2] = e_5, \quad [e_1, e_4] = [e_2, e_3] = e_6, \\ \mathfrak{n}_6 &: [e_1, e_2] = e_5, \quad [e_1, e_4] = [e_2, e_5] = e_6, \\ \mathfrak{n}_7 &: [e_1, e_2] = e_4, \quad [e_1, e_3] = -[e_2, e_4] = e_5, \quad [e_1, e_4] = [e_2, e_3] = e_6. \end{aligned}$$

² Given a 2-step solvable Lie algebra $\mathfrak{g}$ and an abelian ideal $\mathfrak{u}$ of $\mathfrak{g}$ such that $\mathfrak{g}' \subseteq \mathfrak{u}$, the Jacobi identity implies that $\{\text{ad}_x|_{\mathfrak{u}} : x \in \mathfrak{g}\}$ is a commutative family of endomorphisms of $\mathfrak{u}$.

Proof. According to Lemma 2.4, $\mathfrak{n}$ is at most 3-step nilpotent. To carry out the classification, we will consider separately the k -step nilpotent case for $k = 2$ or 3 .

• $\mathfrak{n}$ is 2-step nilpotent: We observe that $\dim \mathfrak{n}' = 1$ or 2 . In fact, the center $\mathfrak{z}$ is a J -stable proper ideal, hence $\dim \mathfrak{z} = 2$ or 4 . If $\dim \mathfrak{z} = 4$, then $\dim \mathfrak{n}' = 1$, while if $\dim \mathfrak{z} = 2$, we have that $\dim \mathfrak{n}' = 1$ or 2 since $\mathfrak{n}' \subset \mathfrak{z}$.

If $\dim \mathfrak{n}' = 1$, according to [4, Theorem 4.1], $\mathfrak{n}$ is isomorphic to $\mathfrak{n}_1$ or $\mathfrak{n}_2$.

If $\dim \mathfrak{n}' = 2$, since $\mathfrak{z}$ is J -stable, then $\mathfrak{n}' = \mathfrak{z}$ and we have that $\dim \mathfrak{z} = 2$. Let $\mathfrak{v}$ be a J -stable complementary subspace of $\mathfrak{z}$. There are two possibilities:

- (i) there exists $x \in \mathfrak{v}$ such that $\text{rank } \text{ad}_x = 1$;
- (ii) $\text{rank } \text{ad}_x = 2$ for any $x \in \mathfrak{v}$, $x \neq 0$.

We consider next each case separately.

– Case (i): Consider the 3-dimensional subspace $W := \ker(\text{ad}_x|_{\mathfrak{v}})$. It follows immediately that $\dim(W \cap JW) = 2$.

If $x \notin W \cap JW$, there exists $e_1 \in W \cap JW$ such that $\{e_1, Je_1, x, Jx\}$ is a basis of $\mathfrak{v}$. Since J is abelian, it follows that the only non-zero brackets are $[e_1, Je_1]$, $[x, Jx]$, which must be linearly independent. Setting $e_3 := x$, $e_4 := Jx$, $e_5 = [e_1, Je_1]$, $e_6 := [x, Jx]$, we obtain that $\mathfrak{n} \cong \mathfrak{n}_3$ and $J = J_{s,t}$ is given by

$$(8) \quad J_{s,t} e_1 = e_2, \quad J_{s,t} e_3 = e_4, \quad J_{s,t} e_5 = se_5 + te_6,$$

for some $s, t \in \mathbb{R}$, $t \neq 0$, since the center $\mathfrak{z} = \text{span}\{e_5, e_6\}$ is J -stable.

If $x \in W \cap JW$, then $\{x, Jx\}$ is a basis of $W \cap JW$. There exists $e_1 \in W$ such that $Je_1 \notin W$ and therefore, $\{e_1, Je_1, x, Jx\}$ is a basis of $\mathfrak{v}$. Since J is abelian, it follows that the only non-zero brackets are $[e_1, Je_1]$ and $[e_1, Jx] = -[Je_1, x]$, which must be linearly independent. Setting $e_2 := Je_1$, $e_3 := -x$, $e_4 = Jx$, $e_5 := [e_1, Je_1]$, $e_6 := [e_1, Jx]$, we obtain that $\mathfrak{n} \cong \mathfrak{n}_5$ and $J = J_{s,t}$ is given by

$$(9) \quad J_{s,t} e_1 = e_2, \quad J_{s,t} e_3 = -e_4, \quad J_{s,t} e_5 = se_5 + te_6,$$

for some $s, t \in \mathbb{R}$ with $t \neq 0$, since the center $\mathfrak{z} = \text{span}\{e_5, e_6\}$ is J -stable.

– Case (ii): Two non-equivalent families of abelian complex structures will appear according to the following possibilities:

- (J^1) there exists $e_1 \in \mathfrak{v}$ such that $W := \ker(\text{ad}_{e_1}|_{\mathfrak{v}})$ is not J -stable,
- (J^2) for any $x \in \mathfrak{v}$, $\ker(\text{ad}_x|_{\mathfrak{v}})$ is J -stable.

(J^1) In this case $\mathfrak{v} = W \oplus JW$ and we fix a basis $\{e_1, y, Je_1, Jy\}$ of $\mathfrak{v}$ with $y \in W$. It follows that $[e_1, y] = 0 = [Je_1, Jy]$. A basis of $\mathfrak{z} = \mathfrak{n}'$ is given by $\{[e_1, Je_1], [e_1, Jy]\}$ since $\text{rank } \text{ad}_{e_1} = 2$. Moreover, $[e_1, Jy] = [y, Je_1]$ since J is abelian and $\text{rank } \text{ad}_y = 2$ implies $[y, Jy] \neq 0$. Therefore, there exist $a, b \in \mathbb{R}$, $a^2 + b^2 \neq 0$ such that

$$[y, Jy] = a[e_1, Je_1] + b[e_1, Jy].$$

Since $\text{rank } \text{ad}_w = 2$ for all $w \in W$, $w \neq 0$, it follows that $\det(\text{ad}_{ce_1+y}|_{JW}) \neq 0$ for any c . We have that $\det(\text{ad}_{ce_1+y}|_{JW}) = c^2 + cb - a$, with respect to the bases $\{Je_1, Jy\}$, $\{[e_1, Je_1], [e_1, Jy]\}$ of JW and $\mathfrak{z}$, respectively. Therefore, it must be $b^2 + 4a < 0$, and setting

$$e_2 := \frac{1}{\sqrt{-(b^2 + 4a)}}(-be_1 + 2y), \quad e_3 := Je_1, \quad e_4 := Je_2, \quad e_5 := [e_1, e_3], \quad e_6 := [e_1, e_4],$$

it turns out that

$$[e_1, e_3] = -[e_2, e_4] = e_5, \quad [e_1, e_4] = [e_2, e_3] = e_6.$$

Hence, $\mathfrak{n} \cong \mathfrak{n}_4$ and $J = J_{s,t}^1$ is given by

$$(10) \quad J_{s,t}^1 e_1 = e_3, \quad J_{s,t}^1 e_2 = e_4, \quad J_{s,t}^1 e_5 = se_5 + te_6,$$

for some $s, t \in \mathbb{R}$ with $t \neq 0$, since the center $\mathfrak{z} = \text{span}\{e_5, e_6\}$ is J -stable.

(J^2) Let $e_1 \in \mathfrak{v}$ and consider a basis $\{e_1, Je_1\}$ of $W_1 := \ker(\text{ad}_{e_1}|_{\mathfrak{v}})$. Let $x \in \mathfrak{v}$ such that $x \notin W_1$, then $W_2 := \ker(\text{ad}_x|_{\mathfrak{v}})$ is a J -stable complementary subspace of W_1 in $\mathfrak{v}$, with basis $\{x, Jx\}$. Setting $e_2 := Je_1$, $e_3 := Jx$, $e_4 := x$, it turns out that $\mathfrak{n} \cong \mathfrak{n}_4$ and $J = J_{s,t}^2$ is given by

$$(11) \quad J_{s,t}^2 e_1 = e_2, \quad J_{s,t}^2 e_3 = -e_4, \quad J_{s,t}^2 e_5 = se_5 + te_6,$$

for some $s, t \in \mathbb{R}$ with $t \neq 0$, since the center $\mathfrak{z} = \text{span}\{e_5, e_6\}$ is J -stable.

• $\mathfrak{n}$ is 3-step nilpotent: In this case, $\mathfrak{n}^2 = [\mathfrak{n}, \mathfrak{n}'] \neq 0$ and $\mathfrak{n}^2 \subseteq \mathfrak{z}$. The center $\mathfrak{z}$ being J -stable, it follows that $\mathfrak{n}_J^2 = \mathfrak{n}^2 + J\mathfrak{n}^2 \subseteq \mathfrak{z}$. Since $\dim \mathfrak{z} \neq 4$ (otherwise, $\dim \mathfrak{n}' = 1$) it follows that $\mathfrak{n}_J^2 = \mathfrak{z}$ and $\dim \mathfrak{n}_J^2 = 2$. From Lemma 2.4, $\mathfrak{n}_J^2 \subsetneq \mathfrak{n}'_J$, thus $\dim \mathfrak{n}'_J = 4$. Let V_0 be a J -stable subspace complementary to $\mathfrak{n}'_J$ and fix a basis $\{f_1, Jf_1 = f_2\}$ of V_0 . Using that $\mathfrak{n}'_J$ is abelian (Proposition 3.1), we obtain

$$\mathfrak{n}' = [V_0 \oplus \mathfrak{n}'_J, V_0 \oplus \mathfrak{n}'_J] = [V_0, V_0] + [V_0, \mathfrak{n}'_J] \subseteq [V_0, V_0] + \mathfrak{n}^2,$$

thus $[f_1, f_2] \notin \mathfrak{z}$, hence $J[f_1, f_2] \notin \mathfrak{z}$, which implies that $\mathfrak{n}'_J = V_1 \oplus \mathfrak{n}_J^2$, where V_1 is the subspace spanned by $[f_1, f_2]$ and $J[f_1, f_2]$. Note that $\mathfrak{n} = V_0 \oplus V_1 \oplus \mathfrak{n}_J^2$ and that for any $y \in V_0$, $y \neq 0$, the images of $\text{ad}_y : V_1 \rightarrow \mathfrak{n}_J^2$ coincide with $\mathfrak{n}^2$.

– If $\dim \mathfrak{n}^2 = 1$, we may assume that $[f_2, [f_1, f_2]] \neq 0$. By making a suitable change of basis of V_0 , we obtain furthermore that $[f_1, [f_1, f_2]] = 0$. Setting

$$e_1 := f_1, \quad e_2 := f_2, \quad e_3 := J[f_2, [f_1, f_2]], \quad e_4 := -J[f_1, f_2], \quad e_5 := [f_1, f_2], \quad e_6 := [f_2, [f_1, f_2]],$$

we obtain that $\mathfrak{n} \cong \mathfrak{n}_6$ and J is given by

$$(12) \quad Je_1 = e_2, \quad Je_3 = -e_6, \quad Je_4 = e_5.$$

– If $\dim \mathfrak{n}^2 = 2$ the map $\text{ad}_{f_1} : V_1 \rightarrow \mathfrak{n}_J^2$ is an isomorphism and setting

$$e_1 := f_1, \quad e_2 := f_2, \quad e_3 := J[f_1, f_2], \quad e_4 := [f_1, f_2], \quad e_5 := [f_1, J[f_1, f_2]], \quad e_6 := [f_1, [f_1, f_2]]$$

one has a basis of $\mathfrak{n}_7$. Since J is abelian, it leaves $\mathfrak{z} = \text{span}\{e_5, e_6\}$ stable, hence it is given by $J = J_{s,t}$, with

$$(13) \quad J_{s,t}e_1 = e_2, \quad J_{s,t}e_3 = -e_4, \quad J_{s,t}e_5 = se_5 + te_6$$

for some $s, t \in \mathbb{R}$, $t \neq 0$. □

Remark 3.3. It follows from Theorem 3.2 that if $\mathfrak{n}$ is a 6-dimensional nilpotent non-abelian Lie algebra and J is any abelian complex structure, then

- $\mathfrak{n}'_J \cong \mathbb{R}^2$ if and only if $\mathfrak{n} = \mathfrak{n}_k$, $1 \leq k \leq 5$,
- $\mathfrak{n}'_J \cong \mathbb{R}^4$ if and only if $\mathfrak{n} = \mathfrak{n}_6$ or $\mathfrak{n}_7$.

3.1. Equivalence classes of complex structures. In this subsection we parameterize the equivalence classes of abelian complex structures on the Lie algebras appearing in Theorem 3.2.

Let us begin with $\mathfrak{n}_1$ and $\mathfrak{n}_2$. According to [20] (see also Proposition 2.2) it follows that, up to equivalence, $\mathfrak{n}_1$ has a unique complex structure J , given by

$$Je_1 = e_2, \quad Je_3 = e_4, \quad Je_5 = e_6,$$

whereas $\mathfrak{n}_2$ has two, given by

$$J_{\pm} e_1 = e_2, \quad J_{\pm} e_3 = \pm e_4, \quad J_{\pm} e_5 = e_6.$$

For the Lie algebra $\mathfrak{n}_6$, it follows from the proof of Theorem 3.2 that it has a unique abelian complex structure, given by (12).

We will determine next the equivalence classes of abelian complex structures on the remaining Lie algebras. To achieve this, we consider a nilpotent Lie algebra $\mathfrak{n}$ and the following space:

$$\mathcal{C}_a(\mathfrak{n}) = \{J : \mathfrak{n} \rightarrow \mathfrak{n} : J \text{ is an abelian complex structure on } \mathfrak{n}\},$$

which we suppose non-empty. $\mathcal{C}_a(\mathfrak{n})$ is a closed subset of $GL(\mathfrak{n})$. The group $\text{Aut}(\mathfrak{n})$ acts on $\mathcal{C}_a(\mathfrak{n})$ by conjugation, and the quotient $\mathcal{C}_a(\mathfrak{n})/\text{Aut}(\mathfrak{n})$ parameterizes the equivalence classes of abelian complex structures on $\mathfrak{n}$.

For $\mathfrak{n} = \mathfrak{n}_3, \mathfrak{n}_5$ and $\mathfrak{n}_7$, it follows from equations (8), (9) and (13) in the proof of Theorem 3.2 that $\mathcal{Z} := \{(s, t) \in \mathbb{R}^2 : t \neq 0\}$ can be considered as a subspace of $\mathcal{C}_a(\mathfrak{n})$ such that $\text{Aut}(\mathfrak{n}) \cdot \mathcal{Z} = \mathcal{C}_a(\mathfrak{n})$. The induced topology on $\mathcal{Z}$ coincides with the usual topology from $\mathbb{R}^2$. Therefore, we have a homeomorphism

$$\mathcal{C}_a(\mathfrak{n})/\text{Aut}(\mathfrak{n}) \cong \mathcal{Z}/G,$$

where $G := \{\phi \in \text{Aut}(\mathfrak{n}) : \phi(\mathcal{Z}) = \mathcal{Z}\}$, a closed Lie subgroup of $\text{Aut}(\mathfrak{n})$.

Let J be an abelian complex structure on $\mathfrak{n}$ and $\mathfrak{v}$ a J -stable complementary subspace of the center $\mathfrak{z}$ in $\mathfrak{n}$. It can be shown that $\text{Aut}^0(\mathfrak{n})$ acts transitively on the orbit of J in $\mathcal{C}_a(\mathfrak{n})$, that is,

$$\text{Aut}(\mathfrak{n})J = \text{Aut}^0(\mathfrak{n})J,$$

where $\text{Aut}^0(\mathfrak{n}) = \{\phi \in \text{Aut}(\mathfrak{n}) : \phi(\mathfrak{v}) = \mathfrak{v}\}$. Indeed, if $J' = \phi J \phi^{-1}$ with $\phi = \begin{pmatrix} A & 0 \\ B & C \end{pmatrix} \in \text{Aut}(\mathfrak{n})$, then $\phi^0 = \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix} \in \text{Aut}^0(\mathfrak{n})$ satisfies $J' = \phi^0 J (\phi^0)^{-1}$. This implies that

$$\mathcal{C}_a(\mathfrak{n})/\text{Aut}(\mathfrak{n}) \cong \mathcal{Z}/G^0$$

for the Lie algebras $\mathfrak{n}_i$, $i = 3, 5, 7$, where $G^0 = G \cap \text{Aut}^0(\mathfrak{n})$.

For $\mathfrak{n} = \mathfrak{n}_4$, equations (10) and (11) in the proof of Theorem 3.2 give two non-equivalent families $\{J_{s,t}^1\}, \{J_{s,t}^2\}$, $t \neq 0$, hence we have two associated subspaces $\mathcal{Z}_1, \mathcal{Z}_2$ of $\mathcal{C}_a(\mathfrak{n}_4)$ such that

$$\mathcal{C}_a(\mathfrak{n})/\text{Aut}(\mathfrak{n}) \cong \mathcal{Z}_1/G_1^0 \dot{\cup} \mathcal{Z}_2/G_2^0.$$

We will determine in what follows for each algebra $\mathfrak{n}_i$, $i = 3, 5, 7$, the group G^0 and orbit space $\mathcal{Z}/G^0$ (respectively, G_j^0 and $\mathcal{Z}_j/G_j^0$, $j = 1, 2$, in the case of $\mathfrak{n}_4$).

- $\mathfrak{n}_3$: In this case, the group G^0 is given by a disjoint union $G^0 = G^+ \dot{\cup} G^-$, where

$$G^+ = \left\{ \begin{pmatrix} a & -b & & & \\ b & a & & & \\ & & c & -d & \\ & & d & c & \\ & & & & a^2 + b^2 \\ & & & & & c^2 + d^2 \end{pmatrix} : \begin{matrix} a^2 + b^2 \neq 0, \\ c^2 + d^2 \neq 0 \end{matrix} \right\}, \quad G^- = \begin{pmatrix} & & & I \\ & & & \\ I & & & \\ & & 0 & 1 \\ & & 1 & 0 \end{pmatrix} \cdot G^+.$$

Let $(s, t), (s', t') \in \mathcal{Z}$ such that $\phi \cdot (s, t) = (s', t')$, with $\phi \in G^0$. It can be shown that:

- (i) if $\phi \in G^+$, then $s' = s$ and $tt' > 0$;
- (ii) if $\phi \in G^-$, then $s' = -s$ and $tt' < 0$.

Let us fix now $(s, t) \in \mathcal{Z}$. Taking $\phi_+ \in G^+$ with $a = \sqrt{|t|}$, $c = 1$ and $b = d = 0$, we obtain that $\phi_+ \cdot (s, t) = (s, \frac{t}{|t|})$. Taking now $\phi_- \in G^-$ with $a = 1$, $c = \sqrt{1 + s^2}$ and $b = d = 0$, we obtain that $\phi_- \cdot (s, -1) = (-s, 1)$. To sum up, we have that

$$J_{s,t} \text{ is equivalent to either } \begin{cases} J_{s,1} & \text{if } t > 0, \\ J_{-s,1} & \text{if } t < 0. \end{cases}$$

It follows from (i) and (ii) that $J_{s,1}$ and $J_{s',1}$ are equivalent if and only if $s = s'$, hence

$$\mathcal{C}_a(\mathfrak{n}_3) / \text{Aut}(\mathfrak{n}_3) \cong \{J_{s,1} : s \in \mathbb{R}\}.$$

Remark 3.4. In [14] a curve of non-equivalent abelian complex structures was obtained by using geometric invariant theory.

- $\mathfrak{n}_5$: In this case, the action of G on $\mathcal{Z}$ is transitive, since the automorphism ϕ of $\mathfrak{n}_5$ given by

$$\phi = \begin{pmatrix} 0 & -1 & & & \\ 1 & 0 & & & \\ 0 & -\frac{s}{2} & 0 & \frac{1+s^2}{t} & \\ -\frac{s}{2} & 0 & -\frac{1+s^2}{t} & 0 & \\ & & & 1 & 0 \\ & & & -s & \frac{1+s^2}{t} \end{pmatrix}$$

satisfies $\phi \cdot (s, t) = (0, 1)$ for any $(s, t) \in \mathcal{Z}$. Thus, $\mathcal{C}_a(\mathfrak{n}_5) / \text{Aut}(\mathfrak{n}_5) = \{J_{(0,1)}\}$.

- $\mathfrak{n}_7$: In this case the subgroup G^0 of G is given by

$$\mathbb{C}^\times = \left\{ \begin{pmatrix} a & -b & & & \\ b & a & & & \\ & & d & & \\ & & & d & \\ & & & & ad & -bd \\ & & & & bd & ad \end{pmatrix} : a, b \in \mathbb{R}, d = a^2 + b^2 \neq 0 \right\}.$$

We note that $(0, 1)$ and $(0, -1)$ are fixed points by the action of $\mathbb{C}^\times$. If $(s, t), (s', t') \in \mathcal{Z} - \{(0, \pm 1)\}$, $(s, t) \neq (s', t')$ and $\phi \cdot (s, t) = (s', t')$, with $\phi \in \mathbb{C}^\times$, then by writing down the equations, it can be shown that $tt' > 0$ and both (s, t) and (s', t') lie in the circle

$$(14) \quad S_c := \left\{ (u, v) \in \mathcal{Z} : u^2 + \left(v - \frac{c}{2}\right)^2 = \left(\frac{c}{2}\right)^2 - 1 \right\}, \quad c = t + \frac{1+s^2}{t} = t' + \frac{1+s'^2}{t'}.$$

Each of these circles intersects the v -axis in points $(0, t_0)$ and $(0, 1/t_0)$, with $0 < |t_0| < 1$, where $t_0 = \frac{1}{2}(c \mp \sqrt{c^2 - 1})$, depending on the sign of t . We show next that each S_c coincides with an orbit $\mathcal{O}_{0,t_0}$ of a point under the action of $\mathbb{C}^\times$. Since $\mathbb{C}^\times \cong \mathbb{R}_+ \times S^1$ as Lie groups, and the action of $\mathbb{R}_+$ on $\mathcal{Z}$ is trivial, we only have to consider the action of S^1 on $\mathcal{Z}$. The isotropy group of this action is $\{\pm 1\}$ and therefore each orbit is homeomorphic to $\mathbb{R}P^1 \cong S^1$. Furthermore, each orbit is contained in one circle S_c for some c , hence, by using topological arguments, we have that each S_c coincides with an orbit $\mathcal{O}_{0,t_0}$ (see Figure 3.1), so that,

$$\mathcal{C}_a(\mathfrak{n}_7)/\text{Aut}(\mathfrak{n}_7) \cong \{J_{0,t} : 0 < |t| \leq 1\}.$$

We note that the topology of the quotient space coincides with relative topology of $\mathbb{R}$, in particular, the moduli space of abelian complex structures on $\mathfrak{n}_7$ is disconnected.

• $\mathfrak{n}_4$: The group G_1^0 is given by a disjoint union $G_1^0 = G_1^+ \dot{\cup} G_1^-$, where

$$G_1^+ = \left\{ \begin{pmatrix} a & -\epsilon b & & & \\ b & \epsilon a & & & \\ & & a & -\epsilon b & \\ & & b & \epsilon a & \\ & & & & a^2 - b^2 & -2\epsilon ab \\ & & & & 2ab & \epsilon(a^2 - b^2) \end{pmatrix} : \begin{array}{l} a, b \in \mathbb{R}, \\ a^2 + b^2 \neq 0, \\ \epsilon = \pm 1 \end{array} \right\}, \quad G_1^- = \begin{pmatrix} & -I & & \\ I & & & \\ & & 1 & 0 \\ & & 0 & 1 \end{pmatrix} \cdot G_1^+.$$

We note that the G_1^0 -orbit of $(0, 1)$ has two points: $(0, \pm 1)$. The orbit $\mathcal{O}_{s,t}$ of $(s, t) \in \mathcal{Z}_1 - \{(0, \pm 1)\}$ is given by the disjoint union of two circles (see Figure 3.1):

$$(15) \quad \mathcal{O}_{s,t} := \left\{ (u, v) \in \mathcal{Z}_1 : u^2 + \left(v \pm \frac{c}{2}\right)^2 = \left(\frac{c}{2}\right)^2 - 1 \right\},$$

with $c = t + \frac{1+s^2}{t}$. Each of these orbits intersects the positive v -axis in points $(0, t_0)$ and $(0, 1/t_0)$, with $0 < t_0 < 1$. Thus, $\mathcal{Z}_1/G_1^0$ is parameterized by $(0, 1]$.


In a similar manner, one shows that the group G_2^0 is given by a disjoint union $G_2^0 = G_2^+ \dot{\cup} G_2^-$, where

$$(16) \quad G_2^+ = \left\{ \begin{pmatrix} a & -b & & & \\ b & a & & & \\ & & c & -d & \\ & & d & c & \\ & & & & ac - bd & -(ad + bc) \\ & & & & ad + bc & ac - bd \end{pmatrix} : \begin{array}{l} a, b, c, d \in \mathbb{R}, \\ a^2 + b^2 \neq 0, \\ c^2 + d^2 \neq 0 \end{array} \right\}, \quad G_2^- = G_2^+ \cdot \begin{pmatrix} & U & & \\ & & & \\ U & & & \\ & & & -U \end{pmatrix},$$

with $U = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. We have that $G_2^+ \cong \mathbb{C}^\times \times \mathbb{C}^\times \cong \mathbb{R}_+^2 \times T^2$ with $\mathbb{R}_+^2$ acting trivially on $\mathcal{Z}_2$.

The isotropy of the T^2 -action is the subgroup $\{(z, w) \in S^1 \times S^1 : zw = \pm 1\}$. Therefore, for $(s, t) \in \mathcal{Z}_2 - \{(0, \pm 1)\}$, the G_2^+ -orbit through (s, t) is a circle as in (14). From (16) it follows that the G_2^- -orbit through (s, t) is also a circle and therefore the orbit $\mathcal{O}_{s,t}$ is given as in (15), therefore, $\mathcal{Z}_2/G_2^0 \cong (0, 1]$, giving the following homeomorphism:

$$\mathcal{C}_a(\mathfrak{n}_4)/\text{Aut}(\mathfrak{n}_4) \cong (0, 1] \times \mathbb{Z}_2.$$



The results in this section are summarized in the next theorem.

Theorem 3.5. *Let $\mathfrak{n}$ be a 6-dimensional nilpotent Lie algebra with an abelian complex structure J . Then $(\mathfrak{n}, J)$ is holomorphically isomorphic to one and only one of the following:*

- (1) $(\mathfrak{n}_1, J)$, with its unique complex structure: $Je_1 = e_2$, $Je_3 = e_4$, $Je_5 = e_6$,
- (2) $(\mathfrak{n}_2, J_{\pm})$, with $J_{\pm}e_1 = e_2$, $J_{\pm}e_3 = \pm e_4$, $J_{\pm}e_5 = e_6$,
- (3) $(\mathfrak{n}_3, J_s)$, with $J_s e_1 = e_2$, $J_s e_3 = e_4$, $J_s e_5 = se_5 + e_6$, $s \in \mathbb{R}$,
- (4) $(\mathfrak{n}_4, J_t^1)$ with $J_t^1 e_1 = e_3$, $J_t^1 e_2 = e_4$, $J_t^1 e_5 = te_6$, $t \in (0, 1]$,
- (5) $(\mathfrak{n}_4, J_t^2)$ with $J_t^2 e_1 = e_2$, $J_t^2 e_3 = -e_4$, $J_t^2 e_5 = te_6$, $t \in (0, 1]$,
- (6) $(\mathfrak{n}_5, J)$ with $Je_1 = e_2$, $Je_3 = -e_4$, $Je_5 = e_6$,
- (7) $(\mathfrak{n}_6, J)$, with $Je_1 = e_2$, $Je_3 = -e_6$, $Je_4 = e_5$,
- (8) $(\mathfrak{n}_7, J_t)$ with $J_t e_1 = e_2$, $J_t e_3 = -e_4$, $J_t e_5 = te_6$, $0 < |t| \leq 1$.

Remark 3.6. It was proved in [20] that every complex structure on $\mathfrak{n}_6$ is abelian. This fact, together with Theorem 3.5, implies that $\mathfrak{n}_6$ admits a unique complex structure up to equivalence.

Remark 3.7. We observe that $\mathfrak{n}_4$ has a bi-invariant complex structure given by

$$J_0 e_1 = e_2, \quad J_0 e_3 = e_4, \quad J_0 e_5 = e_6.$$

According to [13] J_0 is the unique bi-invariant complex structure on $\mathfrak{n}_4$ compatible with the standard orientation (see also [1]).

Notation. The Lie algebras in Theorem 3.2 appear in the literature with different notations. In [10], the nilpotent Lie algebras equipped with the so-called nilpotent complex structures were denoted by $\mathfrak{h}_k$, $k = 1, \dots, 16$. Salamon in [18] associates to each Lie algebra a 6-tuple encoding the expression of the differential of 1-forms. We give next the correspondence among the various notations:

$$\begin{aligned}
 \mathfrak{n}_1 &\longleftrightarrow \mathfrak{h}_8 \longleftrightarrow (0, 0, 0, 0, 0, 12), \\
 \mathfrak{n}_2 &\longleftrightarrow \mathfrak{h}_3 \longleftrightarrow (0, 0, 0, 0, 0, 12 + 34), \\
 \mathfrak{n}_3 &\longleftrightarrow \mathfrak{h}_2 \longleftrightarrow (0, 0, 0, 0, 12, 34), \\
 \mathfrak{n}_4 &\longleftrightarrow \mathfrak{h}_5 \longleftrightarrow (0, 0, 0, 0, 13 + 42, 14 + 23), \\
 \mathfrak{n}_5 &\longleftrightarrow \mathfrak{h}_4 \longleftrightarrow (0, 0, 0, 0, 12, 14 + 23), \\
 \mathfrak{n}_6 &\longleftrightarrow \mathfrak{h}_9 \longleftrightarrow (0, 0, 0, 0, 12, 14 + 25), \\
 \mathfrak{n}_7 &\longleftrightarrow \mathfrak{h}_{15} \longleftrightarrow (0, 0, 0, 12, 13 + 42, 14 + 23).
 \end{aligned}$$

4. DIMENSION SIX: THE GENERAL CASE

The classification, up to equivalence, of the 6-dimensional non-nilpotent Lie algebras $\mathfrak{s}$ admitting abelian complex structures J will be done in several steps, according to the different possibilities for $\mathfrak{s}'_J$. When J is proper it follows from §2.3 that the ideal $\mathfrak{s}'_J$ with its induced abelian complex structure coincides, up to equivalence, with $\mathbb{R}^2$, $\mathfrak{aff}(\mathbb{R})$, $\mathbb{R}^4$, $\mathfrak{h}_3 \times \mathbb{R}$, $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$, $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$, $\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2$ endowed with their unique abelian complex structure or with $(\mathfrak{aff}(\mathbb{C}), J_1)$, $(\mathfrak{aff}(\mathbb{C}), J_2)$. Note that according to Proposition 3.1, the case $\mathfrak{s}'_J \cong \mathfrak{h}_3 \times \mathbb{R}$ does not occur. In order to carry out the classification we will consider three different cases: $\dim \mathfrak{s}'_J = 2$, $\dim \mathfrak{s}'_J = 4$ or $\mathfrak{s}'_J = \mathfrak{s}$.

4.1. $\dim \mathfrak{s}'_J = 2$.

Theorem 4.1. *If $\mathfrak{s}$ is a 6-dimensional non-nilpotent Lie algebra admitting an abelian complex structure J such that $\dim \mathfrak{s}'_J = 2$, then $(\mathfrak{s}, J)$ is holomorphically isomorphic to one of the following:*

- $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^4$ with its unique complex structure,
- $\mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ equipped with the product $(J_1 \times J)$, J_1 from Theorem 2.5.

Proof. Since $\dim \mathfrak{s}'_J = 2$, we have two possibilities:

- (1) $\dim \mathfrak{s}' = 1$. Since $\mathfrak{s}$ is non-nilpotent, $\mathfrak{s}$ is isomorphic to $\mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^4$ (see, for instance, [4, Theorem 4.1]). According to Proposition 2.2, this Lie algebra has a unique complex structure, up to equivalence.
- (2) $\dim \mathfrak{s}' = 2$, hence $\mathfrak{s}' = \mathfrak{s}'_J$ is J -stable. Let $\mathfrak{s}'$ be spanned by e_1, e_2 and $Je_1 = e_2$. Let $\mathfrak{u}$ be a complementary J -stable subspace of $\mathfrak{s}$, $\mathfrak{s} = \mathfrak{s}' \oplus \mathfrak{u}$. From Lemma 2.3(iii), the kernel of the linear map $\rho : \mathfrak{u} \rightarrow \text{End}(\mathfrak{s}')$, $u \mapsto \text{ad}_u|_{\mathfrak{s}'}$ is J -stable. The Jacobi identity implies that $\text{Im} \rho$ is a commutative family of endomorphisms of $\mathfrak{s}'$, therefore $\dim(\text{Im} \rho) < 4$, thus $\dim(\ker \rho)$ is 4 or 2. When $\dim(\ker \rho) = 4$, $\mathfrak{s}$ is 2-step nilpotent, contradicting the assumption on $\mathfrak{s}$. Indeed, if $\rho \equiv 0$, then $[x, u] = 0$ for all $x \in \mathfrak{s}'$ and $u \in \mathfrak{u}$. Since $\mathfrak{s}'$ is abelian, we obtain that $\mathfrak{s}'$ coincides with the center of $\mathfrak{s}$ and the claim follows. Therefore, we must have $\dim(\ker \rho) = 2$. We show next that $(\mathfrak{s}, J)$ is holomorphically isomorphic to $\mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ equipped with the product $(J_1 \times J)$ from Theorem 2.5. Indeed, let $f_1, f_2 = Jf_1, f_3, f_4 = Jf_3$ be a basis of $\mathfrak{u}$ with $f_1, f_2 \in \ker \rho$. Using the Jacobi identity one obtains $[[f_1, f_2], f_3] = [[f_1, f_2], Jf_3] = 0$, therefore, since J is abelian, $[f_3, [f_1, f_2]] = [f_3, J[f_1, f_2]] = 0$. Thus, $[f_1, f_2] = 0$, since otherwise $f_3 \in \ker \rho$, which is a contradiction. Since $\rho(f_3)\rho(f_4) = \rho(f_4)\rho(f_3)$ and $\rho(f_4) = -\rho(f_3)J$ one has

$$\rho(f_3) = \begin{pmatrix} x & -y \\ y & x \end{pmatrix},$$

with $x^2 + y^2 \neq 0$. We note that $\mathfrak{k} := \text{span}\{f_3, f_4, e_1, e_2\}$ is a J -stable Lie subalgebra of $\mathfrak{s}$, and using Lemma 2.10 we obtain that $(\mathfrak{k}, J|_{\mathfrak{k}})$ is holomorphically isomorphic to $(\mathfrak{aff}(\mathbb{C}), J_1)$, so that we may assume

$$[f_3, f_4] = [e_1, e_2] = 0, \quad [f_3, e_1] = -[f_4, e_2] = e_1, \quad [f_3, e_2] = [f_4, e_1] = e_2.$$

The remaining brackets are given by

$$\begin{aligned} [f_1, f_3] &= ae_1 + be_2 = -[f_2, f_4], \\ [f_1, f_4] &= -be_1 + ae_2 = [f_2, f_3], \end{aligned}$$

where the last line is a consequence of $[[f_2, f_3], f_4] + [[f_4, f_2], f_3] = 0$ and J abelian. If $a = b = 0$, then $\mathfrak{s} = \mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$. If $a^2 + b^2 \neq 0$, one considers

$$\tilde{f}_1 = e_1 + \frac{1}{a^2 + b^2}(af_1 - bf_2), \quad \tilde{f}_2 = J\tilde{f}_1 = e_2 + \frac{1}{a^2 + b^2}(bf_1 + af_2),$$

and finds that $\tilde{f}_1, \tilde{f}_2$ are central elements in $\mathfrak{s}$. Therefore $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ equipped with the complex structure given in the statement. $\square$

4.2. $\dim \mathfrak{s}'_J = 4$.

Let $\mathfrak{s}$ be a 6-dimensional solvable Lie algebra equipped with an abelian complex structure J such that $\dim \mathfrak{s}'_J = 4$. Let $f_1, f_2 = Jf_1$ span a complementary subspace of $\mathfrak{s}'_J$, and set $D := \text{ad}_{f_1}|_{\mathfrak{s}'_J}$. It then follows from Lemma 2.3(ii) that $\text{ad}_{f_2}|_{\mathfrak{s}'_J} = -DJ$; therefore D and DJ are both derivations of $\mathfrak{s}'_J$. The bracket $[f_1, f_2]$ lies in $\mathfrak{s}' \subseteq \mathfrak{s}'_J$, and using the Jacobi identity for f_1, f_2 and any $x \in \mathfrak{s}'_J$, we obtain that

$$(17) \quad \text{ad}_{[f_1, f_2]}|_{\mathfrak{s}'_J} = DJD - D^2J.$$

In what follows we analyze separately two different cases, corresponding to $\mathfrak{s}'_J$ being abelian or non-abelian.

Theorem 4.2. *Let $\mathfrak{s}$ be a 6-dimensional Lie algebra with an abelian complex structure J such that $\mathfrak{s}'_J$ is 4-dimensional and non-abelian. Then $(\mathfrak{s}, J)$ is holomorphically isomorphic to one of the following:*

- (1) $\mathfrak{aff}(\mathbb{R}) \times (\mathfrak{h}_3 \times \mathbb{R})$, with the unique complex structure on each factor,
- (2) $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{C})$, where the complex structure on $\mathfrak{aff}(\mathbb{C})$ is given by J_1 (Theorem 2.5),
- (3) $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$, where each factor carries its unique complex structure,
- (4) $(\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2) \times \mathbb{R}^2$, where each factor is equipped with its unique abelian complex structure,
- (5) $\mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$, where the first factor is equipped with J_2 from Theorem 2.5.

Proof. We will consider several cases, according to the isomorphism class of $\mathfrak{s}'_J$.

(i) Case $\mathfrak{s}'_J \cong \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$: There exists a basis $\{e_1, \dots, e_4\}$ of $\mathfrak{s}'_J$ such that $[e_1, e_2] = e_2$ and J is given by $Je_1 = e_2, Je_3 = e_4$. A derivation D of $\mathfrak{s}'_J$ such that DJ is also a derivation takes the following form:

$$D = \begin{pmatrix} 0 & 0 & & \\ a & b & & \\ & & c & e \\ & & d & f \end{pmatrix},$$

with $a, b, \dots, f \in \mathbb{R}$. From (17), we obtain that

$$e = -d, \quad f = c, \quad [f_1, f_2] = (a^2 + b^2)e_2 + u, \quad u \in \text{span}\{e_3, e_4\}.$$

Since $\mathfrak{s}'_J = \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$, it follows that $c^2 + d^2 \neq 0$ or $u \neq 0$.

We show next that we may assume $a = b = 0$. Indeed, if $a^2 + b^2 \neq 0$, consider another complementary subspace of $\mathfrak{s}'_J$ spanned by

$$\tilde{f}_1 = e_1 - \frac{bf_1 + af_2}{a^2 + b^2}, \quad \tilde{f}_2 = J\tilde{f}_1 = e_2 + \frac{af_1 - bf_2}{a^2 + b^2}.$$

The only non-vanishing brackets with respect to the basis above are

$$[e_1, e_2] = e_2, [\tilde{f}_1, \tilde{f}_2] = \tilde{u}, [\tilde{f}_1, e_3] = [\tilde{f}_2, e_4] = ce_3 + de_4, [\tilde{f}_1, e_4] = -[\tilde{f}_2, e_3] = -de_3 + ce_4,$$

with $\tilde{u} \in \text{span}\{e_3, e_4\}$.

Note that $\mathfrak{s}$ can be decomposed as $\text{span}\{e_1, e_2\} \oplus \text{span}\{\tilde{f}_1, \tilde{f}_2, e_3, e_4\}$. If $c = d = 0$, then $u \neq 0$ and there is a holomorphic isomorphism between $\text{span}\{\tilde{f}_1, \tilde{f}_2, e_3, e_4\}$ and $\mathfrak{h}_3 \times \mathbb{R}$, so that $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{R}) \times (\mathfrak{h}_3 \times \mathbb{R})$. If $c^2 + d^2 \neq 0$, using Lemma 2.10 we obtain that $\text{span}\{\tilde{f}_1, \tilde{f}_2, e_3, e_4\}$ is holomorphically isomorphic to $(\mathfrak{aff}(\mathbb{C}), J_1)$, so that, in this case, $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{C})$.

(ii) Case $\mathfrak{s}'_J \cong \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$: There exists a basis $\{e_1, \dots, e_4\}$ of $\mathfrak{s}'_J$ such that $[e_1, e_2] = e_2$, $[e_3, e_4] = e_4$ and J is given by $Je_1 = e_2$, $Je_3 = e_4$. A derivation D of $\mathfrak{s}'_J$ is of the form

$$D = \begin{pmatrix} 0 & 0 & 0 & 0 \\ a & b & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & c & d \end{pmatrix}$$

with $a, b, c, d \in \mathbb{R}$, and we observe that DJ is also a derivation. From (17), we obtain that

$$[f_1, f_2] = (a^2 + b^2)e_2 + (c^2 + d^2)e_4.$$

If $a^2 + b^2 = c^2 + d^2 = 0$, we obtain that $\mathfrak{s} = \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$. If $a^2 + b^2 \neq 0$, $c^2 + d^2 = 0$, then setting

$$(18) \quad \tilde{f}_1 = -e_1 + \frac{bf_1 + af_2}{a^2 + b^2}, \quad \tilde{f}_2 = J\tilde{f}_1 = -e_2 + \frac{-af_1 + bf_2}{a^2 + b^2},$$

we obtain that $[\tilde{f}_1, \tilde{f}_2] = 0$ and $\tilde{f}_1, \tilde{f}_2$ commute with $\text{span}\{e_1, \dots, e_4\}$. Therefore, $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^2$ via a holomorphic isomorphism. The proof for $a^2 + b^2 = 0$, $c^2 + d^2 \neq 0$ is analogous. Finally, when $a^2 + b^2 \neq 0$, $c^2 + d^2 \neq 0$, we define $\tilde{f}_1, \tilde{f}_2$ as in (18) and this case reduces to the previous one.

(iii) Case $\mathfrak{s}'_J \cong \mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2$: There exists a basis $\{e_1, \dots, e_4\}$ of $\mathfrak{s}'_J$ such that $[e_1, e_2] = e_2$, $[e_1, e_4] = e_4$, $[e_2, e_3] = e_4$ and $Je_1 = e_2$, $Je_3 = -e_4$. A derivation D of $\mathfrak{s}'_J$ such that DJ is also a derivation is of the form:

$$D = \begin{pmatrix} 0 & 0 & 0 & 0 \\ d & c & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a & b & -d & c \end{pmatrix},$$

with $a, b, c, d \in \mathbb{R}$. Using (17), we obtain that

$$[f_1, f_2] = (c^2 + d^2)e_2 + 2(ad + bc)e_4.$$

If $a^2 + b^2 = c^2 + d^2 = 0$, then $\mathfrak{s} = (\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2) \times \mathbb{R}^2$. If $c^2 + d^2 \neq 0$, $a^2 + b^2 = 0$, then setting

$$(19) \quad \tilde{f}_1 = -e_1 + \frac{cf_1 + df_2}{c^2 + d^2}, \quad \tilde{f}_2 = J\tilde{f}_1 = -e_2 + \frac{-df_1 + cf_2}{c^2 + d^2},$$

one has that $[\tilde{f}_1, \tilde{f}_2] = 0$ and $\tilde{f}_1, \tilde{f}_2$ commute with $\text{span}\{e_1, \dots, e_4\}$, and therefore $\mathfrak{s} \cong (\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2) \times \mathbb{R}^2$ via a holomorphic isomorphism. If $c^2 + d^2 = 0$, $a^2 + b^2 \neq 0$, we have again the holomorphic isomorphism $\mathfrak{s} \cong (\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2) \times \mathbb{R}^2$ by considering the following change of basis:

$$\tilde{f}_1 = f_1 + be_3 + ae_4, \quad \tilde{f}_2 = J\tilde{f}_1 = f_2 + ae_3 - be_4.$$

Finally, when $c^2 + d^2 \neq 0$, $a^2 + b^2 \neq 0$, by applying (19) this case reduces to the previous one.

(iv) Case $\mathfrak{s}'_J \cong \mathfrak{aff}(\mathbb{C})$: There exists a basis $\{e_1, \dots, e_4\}$ of $\mathfrak{s}'_J$ such that

$$[e_1, e_3] = -[e_2, e_4] = e_3, \quad [e_1, e_4] = [e_2, e_3] = e_4.$$

It is known that a derivation D of $\mathfrak{aff}(\mathbb{C})$ has matrix form

$$D = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ c & -d & a & -b \\ d & c & b & a \end{pmatrix}$$

with respect to $e_1, \dots, e_4$, for some $a, b, c, d \in \mathbb{R}$. From Theorem 2.5, we may assume that the restriction of J to $\mathfrak{s}'_J$ is J_1 or J_2 .

Let us consider first the case when the restriction is J_1 . It turns out that DJ_1 is also a derivation of $\mathfrak{s}'_{J_1}$, and using (17), we obtain that

$$[f_1, f_2] = 2(ad + bc)e_3 + 2(bd - ac)e_4,$$

for some $a, b, c, d \in \mathbb{R}$. It follows that $\dim \mathfrak{s}'_J = 2$, and therefore, this case cannot occur.

If the restriction is J_2 , we have that DJ_2 is also a derivation of $\mathfrak{s}'_{J_2}$, and using (17), we obtain that

$$[f_1, f_2] = ((a^2 - b^2) + (c^2 - d^2))e_3 + 2(ab + cd)e_4,$$

with $a, b, c, d \in \mathbb{R}$.

If $a^2 + b^2 = c^2 + d^2 = 0$, then $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$. If $c^2 + d^2 \neq 0$, $a^2 + b^2 = 0$, then setting

$$(20) \quad \tilde{f}_1 = f_1 + ce_3 + de_4, \quad \tilde{f}_2 = Jf_1 = f_2 - ce_1 - de_2,$$

we obtain that $[\tilde{f}_1, \tilde{f}_2] = 0$ and $\tilde{f}_1, \tilde{f}_2$ commute with $\text{span}\{e_1, \dots, e_4\}$. Therefore, $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ via a holomorphic isomorphism. The proof for $c^2 + d^2 = 0$, $a^2 + b^2 \neq 0$ is analogous. Finally, when $c^2 + d^2 \neq 0$, $a^2 + b^2 \neq 0$, we define $\tilde{f}_1, \tilde{f}_2$ as in (20) and this case reduces to the previous one. $\square$

We prove next some properties of $(\mathfrak{s}, J)$ for a 6-dimensional non-nilpotent Lie algebra $\mathfrak{s}$ with an abelian complex structure J such that $\mathfrak{s}'_J \cong \mathbb{R}^4$. Given $x \notin \mathfrak{s}'_J$, we denote by $D_x : \mathfrak{s}'_J \rightarrow \mathfrak{s}'_J$ the restriction of ad_x to $\mathfrak{s}'_J$.

Lemma 4.3. *Let $\mathfrak{s}$ be a 6-dimensional non-nilpotent Lie algebra with center $\mathfrak{z}$ (possibly trivial) and denote $\mathfrak{s}^2 = [\mathfrak{s}, \mathfrak{s}']$. If J is an abelian complex structure on $\mathfrak{s}$ such that $\mathfrak{s}'_J \cong \mathbb{R}^4$, then*

- (i) D_x commutes with $D_x J$ for any $x \notin \mathfrak{s}'_J$.
- (ii) $\text{Im} D_x = \mathfrak{s}^2$ for any $x \notin \mathfrak{s}'_J$.
- (iii) $\dim \mathfrak{s}^2 = 2$ or 4 .
- (iv) $\ker D_x = \mathfrak{z}$ for any $x \notin \mathfrak{s}'_J$.
- (v) $\mathfrak{s}'_J = \mathfrak{s}^2 \oplus \mathfrak{z}$.
- (vi) There exists $x \notin \mathfrak{s}'_J$ such that $[x, Jx] \in \mathfrak{z}$.

Proof. (i) Since J is abelian, $D_{Jx} = -D_x J$. Using that $\mathfrak{s}'_J$ is abelian and the Jacobi identity (see (17)), (i) follows.

(ii) Let $x \notin \mathfrak{s}'_J$ and set $V := \text{span}\{x, Jx\}$. We have that $\mathfrak{s} = V \oplus \mathfrak{s}'_J$, therefore

$$\mathfrak{s}^2 = [\mathfrak{s}, \mathfrak{s}'] = [V, \mathfrak{s}'] = D_x \mathfrak{s}' + D_{Jx} \mathfrak{s}' = D_x \mathfrak{s}' + D_x J \mathfrak{s}' = D_x \mathfrak{s}'_J.$$

(iii) Let $x \notin \mathfrak{s}'_J$, then it follows from (ii) that (iii) is equivalent to $\dim(\operatorname{Im} D_x) = 2$ or 4 . We note first that $\operatorname{Im} D_x = \{0\}$ is not possible because this would give $\dim \mathfrak{s}'_J \leq 2$.

If $\dim(\operatorname{Im} D_x) = 1$, let $\operatorname{Im} D_x = \mathbb{R}y$. There exist $a, b \in \mathbb{R}$ such that $D_x y = ay$, $D_x Jy = by$. From (i) $D_x^2 = -(D_x J)^2$, and this implies $a = b = 0$. Therefore, $D_x^2 = D_{Jx}^2 = 0$ and $\mathfrak{s}$ is nilpotent, which is not possible since $\mathfrak{s}$ is non-nilpotent by assumption.

If $\dim(\operatorname{Im} D_x) = 3$, then $\dim(\ker D_x) = 1$, that is, $\ker D_x = \mathbb{R}y$. From (i) we have $D_x(D_x Jy) = D_x J(D_x y) = 0$, hence $D_x Jy = cy$. Using again that $D_x^2 = -(D_x J)^2$, we obtain $0 = D_x^2 y = -(D_x J)^2 y = -c^2 y$, hence $c = 0$. This implies that $Jy \in \ker D_x = \mathbb{R}y$, a contradiction, and (iii) follows.

(iv) It follows from (ii) and (iii) that $\mathfrak{z} \subset \mathfrak{s}'_J$. Moreover,

$$(21) \quad \mathfrak{z} = \ker D_x \cap J(\ker D_x), \quad \text{for any } x \notin \mathfrak{s}'_J.$$

If $\dim \mathfrak{s}^2 = 4$, then using (ii) we obtain that $\ker D_x = \{0\}$ for any $x \notin \mathfrak{s}'_J$, hence $\mathfrak{z} = \{0\}$.

If $\dim \mathfrak{s}^2 = 2$, then $\dim(\ker D_x) = 2$ for any $x \notin \mathfrak{s}'_J$. Let us assume that there exists $x \notin \mathfrak{s}'_J$ such that $\ker D_x$ is not J -stable, therefore, $\mathfrak{s}'_J = \ker D_x \oplus J(\ker D_x)$. Using (ii) we obtain that $D_x : J(\ker D_x) \rightarrow \mathfrak{s}^2$ is an isomorphism, then

$$D_x(\mathfrak{s}^2) = D_x D_x J(\ker D_x) = D_x J D_x(\ker D_x) = 0.$$

This implies $\mathfrak{s}^2 = \ker D_x$, which means that $D_x^2 = 0 = D_{Jx}^2$, contradicting the fact that $\mathfrak{s}$ is not nilpotent. Hence, $\ker D_x$ is J -stable for any $x \notin \mathfrak{s}'_J$ and (21) implies (iv).

(v) If $\dim \mathfrak{s}^2 = 4$ there is nothing to prove (see proof of (iv)). We consider next the case $\dim \mathfrak{s}^2 = 2$, which implies $\dim \mathfrak{z} = 2$. We only need to show that $\mathfrak{s}^2 \cap \mathfrak{z} = \{0\}$. If $\mathfrak{s}^2 \cap \mathfrak{z} \neq \{0\}$, we fix $x \notin \mathfrak{s}'_J$ and consider a J -stable complementary subspace V of $\ker D_x = \mathfrak{z}$ in $\mathfrak{s}'_J$. Using (ii) we obtain that $D_x : V \rightarrow \mathfrak{s}^2$ is an isomorphism. Let $v \in V$, $v \neq 0$, such that $D_x v \in \mathfrak{s}^2 \cap \mathfrak{z}$, then, using (i) and the fact that $\mathfrak{z}$ is J -stable, we have

$$D_x D_x Jv = D_x J D_x v = 0,$$

that is, $D_x Jv \in \ker D_x = \mathfrak{z}$, thus $D_x(V) = \mathfrak{z}$. Therefore, $\mathfrak{s}^2 = \mathfrak{z}$ and $\mathfrak{s}$ is nilpotent, a contradiction, and (v) follows.

(vi) Let $y \notin \mathfrak{s}'_J$, using (v) we have $[y, Jy] = D_y v + w$ with $w \in \mathfrak{z}$. Setting $x = y + \frac{1}{2}Jv$, we compute

$$[x, Jx] = [y, Jy] - \frac{1}{2}D_y v - \frac{1}{2}D_{Jy} Jv = D_y v + w - D_y v = w \in \mathfrak{z},$$

which implies (vi). □

Theorem 4.4. *Let $\mathfrak{s}$ be a 6-dimensional non-nilpotent Lie algebra with an abelian complex structure J such that $\mathfrak{s}'_J \cong \mathbb{R}^4$. Then $(\mathfrak{s}, J)$ is holomorphically isomorphic to one and only one of the following:*

- (1) $(\mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2, J) : [f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$
 $Jf_1 = f_2, J e_1 = -e_2 + e_3, J e_2 = e_1 + e_4, J e_3 = e_4.$
- (2) $(\mathfrak{s}_1, J_1) : [f_1, f_2] = e_3, [f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$
 $J_1 f_1 = f_2, J_1 e_1 = -e_2 + e_3, J_1 e_2 = e_1 + e_4, J_1 e_3 = e_4.$
- (3) $(\mathfrak{s}_1, J_2) : [f_1, f_2] = e_3, \quad [f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$
 $J_2 f_1 = f_2, J_2 e_1 = e_2, J_2 e_3 = e_4,$

- (4) $(\mathfrak{s}_2, J) : [f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$
 $[f_1, e_3] = [f_2, e_4] = e_1 + e_3, \quad [f_1, e_4] = -[f_2, e_3] = e_2 + e_4,$
 $Jf_1 = f_2, \quad Je_1 = e_2, \quad Je_3 = e_4,$
- (5) $(\mathfrak{s}_{(a,b)}, J), (a, b) \in \mathcal{R}$ (see (25)) :
 $[f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$
 $[f_1, e_3] = [f_2, e_4] = ae_3 + be_4, \quad [f_1, e_4] = -[f_2, e_3] = -be_3 + ae_4,$
 $Jf_1 = f_2, \quad Je_1 = e_2, \quad Je_3 = e_4.$

Proof. Let $f_1 \notin \mathfrak{s}'_J$ such that $[f_1, Jf_1] \in \mathfrak{z}$ and denote $f_2 = Jf_1$ (see Lemma 4.3(vi)), $D := D_{f_1}$.

We consider next the different cases according to $\dim \mathfrak{s}^2$, which can be 2 or 4 (see Lemma 4.3(iii)).

• If $\dim \mathfrak{s}^2 = 2$, then there are two possibilities, depending on $\mathfrak{s}^2$ being J -stable or not.

(i) Case $\mathfrak{s}^2$ is J -stable: we note that $[f_1, f_2] \neq 0$, since $\dim \mathfrak{s}'_J = 4$. Let $e_1, e_2 = Je_1$ be a basis of $\mathfrak{s}^2$ and $e_3 = [f_1, f_2]$, $e_4 = Je_3$ a basis of $\mathfrak{z}$. We have

$$D = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}, \quad -DJ = \begin{pmatrix} -AJ_0 & 0 \\ 0 & 0 \end{pmatrix},$$

where $J_0 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and A is a 2×2 non-singular matrix satisfying $AJ_0 = J_0A$, therefore,

$A = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$ with $a^2 + b^2 \neq 0$. Setting

$$(22) \quad \tilde{f}_1 = \frac{1}{\sqrt{a^2 + b^2}}(af_1 + bf_2), \quad \tilde{f}_2 = J\tilde{f}_1 = \frac{1}{\sqrt{a^2 + b^2}}(-bf_1 + af_2),$$

we may assume $A = I$ (observe that $[\tilde{f}_1, \tilde{f}_2] = [f_1, f_2]$), hence the Lie bracket becomes:

$$[f_1, f_2] = e_3, \quad [f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$$

which is the Lie algebra $\mathfrak{s}_1$ in the statement with the complex structure J_2 .

(ii) Case $\mathfrak{s}^2$ is not J -stable: fix $e_1 \in \mathfrak{s}^2$, $e_1 \neq 0$ and set $e_2 \in \mathfrak{s}^2$ such that $Je_1 + e_2 \in \mathfrak{z}$. It can be shown that $\{e_1, e_2\}$ is a basis of $\mathfrak{s}^2$, using that $\mathfrak{z}$ is J -stable. Let $\{e_3, e_4 = Je_3\}$ be a basis of $\mathfrak{z}$; in the basis $\{e_1, \dots, e_4\}$ of $\mathfrak{s}'_J$, D and the restriction of J to $\mathfrak{s}'_J$ are given by

$$D = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}, \quad J = \begin{pmatrix} -J_0 & 0 \\ Y & J_0 \end{pmatrix},$$

with J_0 as above, and $Y \neq 0$ satisfies $YJ_0 = J_0Y$, which implies that Y is non-singular. Also A is non-singular and commutes with J_0 , and using (22), we may assume that $A = I$. By a suitable change of basis in $\mathfrak{z}$, Y can be taken equal to the identity I and J is then given by

$$(23) \quad Jf_1 = f_2, \quad Je_1 = -e_2 + e_3, \quad Je_2 = e_1 + e_4, \quad Je_3 = e_4.$$

If $[f_1, f_2] = 0$, the Lie brackets of $\mathfrak{s}$ are given by

$$[f_1, e_1] = [f_2, e_2] = e_1, \quad [f_1, e_2] = -[f_2, e_1] = e_2,$$

and J is given as in (23). Therefore, $\mathfrak{s} \cong \mathfrak{aff}(\mathbb{C}) \times \mathbb{R}^2$ with a complex structure that preserves the ideal $\mathbb{R}^2$ but does not preserve the ideal $\mathfrak{aff}(\mathbb{C})$.

If $[f_1, f_2] = ae_3 + be_4 \in \mathfrak{z}$, with $a^2 + b^2 \neq 0$, the Lie algebra obtained is isomorphic to $\mathfrak{s}_1$ from (i) above. Indeed, $\phi : \mathfrak{s}_1 \rightarrow \mathfrak{s}$ given by

$$\phi = \begin{pmatrix} 1 & 0 & & & \\ 0 & 1 & & & \\ & & a & -b & \\ & & b & a & \\ & & & & a & -b \\ & & & & b & a \end{pmatrix}$$

is a Lie algebra isomorphism. The complex structure J_1 on $\mathfrak{s}_1$ induced by ϕ is given as in (23). We note that this complex structure is not equivalent to J_2 in (i) since J_2 preserves $\mathfrak{s}^2$ while J_1 does not.

• If $\dim \mathfrak{s}^2 = 4$, then $\mathfrak{z} = 0$, hence $[f_1, f_2] = 0$, D is an isomorphism, and thus D commutes with J . Therefore, we obtain $\mathfrak{s} = \mathbb{R}^2 \ltimes \mathbb{R}^4$, where the complex structure on $\mathbb{R}^4 = \text{span}\{e_1, \dots, e_4\}$ is given by $Je_1 = e_2$, $Je_3 = e_4$. Considering D as an element of $GL(2, \mathbb{C})$, the possible normal Jordan forms for D are

$$\begin{pmatrix} \alpha & 1 \\ 0 & \alpha \end{pmatrix}, \quad \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix},$$

with $\alpha, \beta \in \mathbb{C} - \{0\}$, $|\alpha| \geq |\beta|$. Identifying $\mathbb{R}^2 = \text{span}\{f_1, f_2\}$ with $\mathbb{C} = \text{span}_{\mathbb{C}}\{f_1\}$, we may take $\alpha^{-1}f_1$ and therefore we may assume $\alpha = 1$, $0 < |\beta| \leq 1$. Thinking of D now as an element of $GL(4, \mathbb{R})$, this gives rise to the following possibilities for D , up to complex conjugation:

$$(24) \quad D' = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ & & 1 & 0 \\ & & 0 & 1 \end{pmatrix}, \quad \text{or} \quad D_{(a,b)} = \begin{pmatrix} 1 & 0 & & \\ 0 & 1 & & \\ & & a & -b \\ & & b & a \end{pmatrix}, \quad a, b \in \mathbb{R}, \quad 0 < a^2 + b^2 \leq 1.$$

Let us denote by $\mathfrak{s}_2$ (resp. $\mathfrak{s}_{(a,b)}$) the Lie algebra determined by D' (resp. $D_{(a,b)}$). Clearly, $\mathfrak{s}_2$ is not isomorphic to $\mathfrak{s}_{(a,b)}$ for any (a, b) . Furthermore, it can be shown that the isomorphism classes of the Lie algebras $\mathfrak{s}_{(a,b)}$ are parameterized by

$$(25) \quad \mathcal{R} = \{(a, b) \in \mathbb{R}^2 : 0 < a^2 + b^2 < 1\} \cup \{(a, b) \in S^1 : b \geq 0\}.$$

□

Remark 4.5. We point out that the Lie algebra $\mathfrak{s}_{(a,b)}$ from Theorem 4.4 has a bi-invariant complex structure given by $Jf_1 = -f_2$, $Je_1 = e_2$, $Je_3 = e_4$. Indeed, $\mathfrak{s}_{(a,b)}$ is the realification of the complex Lie algebra $\mathfrak{r}_{3,\lambda}$ with $\lambda = a + ib$, where $\mathfrak{r}_{3,\lambda}$ has a $\mathbb{C}$ -basis $\{w_1, w_2, w_3\}$ with Lie brackets

$$[w_1, w_2] = w_2, \quad [w_1, w_3] = \lambda w_3.$$

The analogous statement holds for $\mathfrak{s}_2$, which is the realification of the complex Lie algebra $\mathfrak{r}_3$ with $\mathbb{C}$ -basis $\{w_1, w_2, w_3\}$ and Lie brackets

$$[w_1, w_2] = w_2, \quad [w_1, w_3] = w_2 + w_3.$$

4.3. $\mathfrak{s}'_J = \mathfrak{s}$. A family of Lie algebras with abelian complex structure can be constructed in the following way. Let A be a commutative associative algebra and set $\mathfrak{aff}(A) := A \oplus A$ with the following Lie bracket and standard complex structure J :

$$(26) \quad [(x, y), (x', y')] = (0, xy' - x'y), \quad J(x, y) = (-y, x),$$

then J is abelian (see [5]). Moreover, J is non-proper if and only if $A^2 = A$.

We show next that any 6-dimensional Lie algebra with a non-proper abelian complex structure is obtained using this procedure.

Theorem 4.6. *Let $\mathfrak{s}$ be a 6-dimensional Lie algebra with a non-proper abelian complex structure J . Then*

- (i) $\dim \mathfrak{s}' = 3$ and $(\mathfrak{s}, J)$ is holomorphically isomorphic to $\mathfrak{aff}(A)$ with its standard complex structure, where A is a 3-dimensional commutative associative algebra such that $A^2 = A$;
- (ii) $(\mathfrak{s}, J)$ is holomorphically isomorphic to one and only one of the following:
 - (1) $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$, where each factor carries its unique complex structure,
 - (2) $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{C})$, where the second factor is equipped with J_2 from Theorem 2.5,
 - (3) $\mathfrak{aff}(\mathbb{R}) \times (\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2)$, where each factor is equipped with its unique abelian complex structure,
 - (4) $(\mathfrak{s}_3, J) : [e_1, f_i] = f_i, i = 1, 2, 3, \quad [e_2, f_1] = f_2, \quad [e_2, f_2] = f_3, \quad [e_3, f_1] = f_3,$
 $Je_i = f_i, i = 1, 2, 3,$
 - (5) $(\mathfrak{s}_4, J) : [e_1, f_i] = f_i, i = 1, 2, 3, \quad [e_2, f_1] = f_2, \quad [e_3, f_1] = f_3,$
 $Je_i = f_i, i = 1, 2, 3.$

Proof. (i) Since $\dim \mathfrak{s} = 6$ and $\mathfrak{s}'_J = \mathfrak{s}$, it follows from (O2) in §2.1 that $\dim \mathfrak{s}' = 3$ or 4. We show first that $\dim \mathfrak{s}' = 4$ is not possible. If $\dim \mathfrak{s}' = 4$ then $\dim(\mathfrak{s}' \cap J\mathfrak{s}') = 2$. We note that J abelian implies $\mathfrak{s}' \cap J\mathfrak{s}' \subset \mathfrak{z}$ and $\mathfrak{s}$ has a basis of the form $\{e_1, e_2, e_3, Je_1, Je_2, Je_3\}$ with $e_1, e_2 \in \mathfrak{s}', e_3 \in \mathfrak{s}' \cap J\mathfrak{s}'$, therefore $\mathfrak{s}' = \text{span}\{[e_1, Je_1], [e_2, Je_2], [e_1, Je_2] = [e_2, Je_1]\}$, contradicting that $\dim \mathfrak{s}' = 4$.

Thus $\dim \mathfrak{s}' = 3$ and $\mathfrak{s} = \mathfrak{s}' \oplus J\mathfrak{s}'$. Note that

$$(27) \quad \mathfrak{s}' = [\mathfrak{s}', J\mathfrak{s}'].$$

The Lie bracket on $\mathfrak{s}$ induces a structure of commutative associative algebra on $\mathfrak{s}'$ in the following way

$$x \cdot y = [x, Jy], \quad \text{for } x, y \in \mathfrak{s}'.$$

Setting $A := (\mathfrak{s}', \cdot)$ it turns out that

$$\phi : \mathfrak{aff}(A) \rightarrow \mathfrak{s}, \quad \phi(x, y) = y - Jx,$$

is a holomorphic isomorphism, where $\mathfrak{aff}(A)$ is equipped with its standard complex structure (26). Moreover, (27) implies that $A^2 = A$ and (i) follows.

(ii) According to (i), in order to obtain the classification of the non-proper abelian complex structures in dimension 6, we must consider $\mathfrak{aff}(A)$ with its standard complex structure, where A is a 3-dimensional commutative associative algebra satisfying $A^2 = A$. It was shown in [12] that there are five isomorphism classes of such associative algebras. These algebras can be realized as matrix algebras as follows:

$$A_1 = \left\{ \begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right\}, \quad A_2 = \left\{ \begin{pmatrix} a & & \\ & b & -c \\ & c & b \end{pmatrix} \right\}, \quad A_3 = \left\{ \begin{pmatrix} a & & \\ & b & c \\ & & b \end{pmatrix} \right\},$$

$$A_4 = \left\{ \begin{pmatrix} a & b & c \\ & a & b \\ & & a \end{pmatrix} \right\}, \quad A_5 = \left\{ \begin{pmatrix} a & 0 & c \\ & a & b \\ & & a \end{pmatrix} \right\},$$

with $a, b, c \in \mathbb{R}$. The Lie brackets on $\mathfrak{aff}(A_i)$, $i = 1, \dots, 5$, together with its standard complex structure are those given in the statement. It is easily verified that these Lie algebras are pairwise non-isomorphic, and this completes the proof of the theorem. $\square$

Remark 4.7. It is straightforward from Theorems 4.1, 4.2 and 4.4 that among the non-nilpotent 6-dimensional Lie algebras carrying proper abelian complex structures, $(\mathfrak{aff}(\mathbb{R}) \ltimes_{\text{ad}} \mathbb{R}^2) \times \mathbb{R}^2$ from Theorem 4.2 is the unique Lie algebra whose nilradical is non-abelian. Indeed, its nilradical is $\mathfrak{h}_3 \times \mathbb{R}^2$. In the non-proper case, only $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$ and $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{aff}(\mathbb{C})$ have abelian nilradical (see Theorem 4.6).

Corollary 4.8. *Let $\mathfrak{s}$ be a 6-dimensional unimodular solvable Lie algebra equipped with an abelian complex structure. Then $\mathfrak{s}$ is isomorphic to $\mathfrak{n}_k$, $k = 1, \dots, 7$, or to $\mathfrak{s}_{(-1,0)}$. Moreover, the simply connected Lie groups corresponding to these Lie algebras admit compact quotients.*

Proof. The first assertion follows easily from Theorems 3.2, 4.1, 4.2, 4.4 and 4.6. The simply connected Lie groups associated to $\mathfrak{n}_k$, $k = 1, \dots, 7$, admit compact quotients since their structure constants are rational (see [15]). The existence of compact quotients of the simply connected solvable Lie group with Lie algebra $\mathfrak{s}_{(-1,0)}$ was proved in [21]. $\square$

REFERENCES

- [1] E. Abbena, S. Garbiero and S. Salamon, *Hermitian geometry on the Iwasawa manifold*, Boll. Un. Mat. Ital. bf 11-B (1997), 231–249.
- [2] M. L. Barberis, *Hypercomplex structures on four dimensional Lie groups*, Proc. Amer. Math. Soc. **125** (4) (1997), 1043–1054.
- [3] M. L. Barberis, I. G. Dotti and R. J. Miatello, *On certain locally homogeneous Clifford manifolds*, Ann. Glob. Anal. Geom. **13** (1995), 289–301.
- [4] M. L. Barberis and I. Dotti, *Hypercomplex structures on a class of solvable Lie groups*, Quart. J. Math. Oxford (2), **47** (1996), 389–404.
- [5] M. L. Barberis and I. Dotti, *Abelian complex structures on solvable Lie algebras*, J. Lie Theory **14** (2004), 25–34.
- [6] M. L. Barberis, I. Dotti and M. Verbitsky, *Canonical bundles of complex nilmanifolds, with applications to hypercomplex geometry*, Math. Research Letters **16**(2) (2009), 331–347, arXiv: 0712.3863.
- [7] T. Christodoulakis, G. O. Papadopoulos and A. Dimakis, *Automorphisms of real 4-dimensional Lie algebras and the invariant characterization of homogeneous 4-spaces*, J. Phys. A: Mathematical & General **36** (2003), 427–442.
- [8] S. Console and A. Fino, *Dolbeault cohomology of compact nilmanifolds*, Transformation Groups **6**(2) (2001), 111–124.
- [9] S. Console, A. Fino and Y. S. Poon, *Stability of abelian complex structures*, Internat. J. Math. **17**(4) (2006), 401–416.
- [10] L. A. Cordero, M. Fernández, A. Gray and L. Ugarte, *Nilpotent complex structures on compact nilmanifolds*, Rend. Circolo Mat. Palermo **49** (Suppl.) (1997), 83–100.
- [11] L. A. Cordero, M. Fernández and L. Ugarte, *Abelian complex structures on 6-dimensional compact nilmanifolds*, Comment. Math. Univ. Carolinae **43**(2) (2002), 215–229.
- [12] M. Goze and E. Remm, *Affine structures on abelian Lie groups*, Linear Algebra Appl. **360** (2003), 215–230.

- [13] G. Ketsetzis and S. Salamon, *Complex structures on the Iwasawa manifold*, Adv. Geom. **4** (2004), 165–179.
- [14] J. Lauret, *Minimal metrics on nilmanifolds*, in Differential geometry and its applications, Matfyzpress, Prague, 2005, 79–97.
- [15] A. Mal'čev, *On a class of homogeneous spaces*, reprinted in Amer. Math. Soc. Trans. Ser. 1 **9** (1962), 276–307.
- [16] A. P. Petravchuk, *Lie algebras decomposable into a sum of an abelian and a nilpotent subalgebra*, Ukr. Math. J. **40** (3) (1988), 331–334.
- [17] S. Rollenske, *Geometry of nilmanifolds with left-invariant complex structure and deformations in the large*, preprint 2009, arXiv:0901.3120, to appear in Proc. London Math. Soc.
- [18] S. Salamon, *Complex structures on nilpotent Lie algebras*, J. Pure Appl. Algebra **157** (2001), 311–333.
- [19] J. E. Snow, *Invariant complex structures on four dimensional solvable real Lie groups*, Manuscripta Math. **66**, (1990), 397–412.
- [20] L. Ugarte, *Hermitian structures on six-dimensional nilmanifolds*, Transformation Groups **12** (1), (2007), 175–202.
- [21] T. Yamada, *A pseudo-Kähler structure on a nontoral compact complex parallelizable solvmanifold*, Geom. Dedicata **112** (2005), 115–122.

C

E-mail address: andrada@famaf.unc.edu.ar

E-mail address: barberis@famaf.unc.edu.ar

E-mail address: idotti@famaf.unc.edu.ar